



2019 Cornell Guide for Integrated Field Crop Management

Cornell Cooperative Extension

These guidelines are not a substitute for pesticide labeling. Always read and understand the product label before using any pesticide.

2019 Cornell Guide for Integrated Field Crop Management

Authors

Jenn Thomas-Murphy (Soil and Crop Sciences Section; *Contact editor*)
Bill Cox (retired) (Soil and Crop Sciences Section; *grain crop production/silage corn hybrids*)
Gary Bergstrom (Plant Pathology and Plant-Microbe Biology Section; *disease management*)
Jerry Cherney (Soil and Crop Sciences Section; *forage production/variety selection*)
Karl Czymmek (Department of Animal Science; *soil fertility/nutrient management*)
Russell Hahn (retired) (Soil and Crop Sciences Section; *weed management*)
Julie Hansen (Plant Breeding and Genetics Section; *forage production/variety selection*)
Michael Helms (Pesticide Management Education Program; *pesticide information*)
Quirine Ketterings (Department of Animal Science; *soil fertility/nutrient management*)
Murray McBride (Soil and Crop Sciences Section; *sewage sludge utilization*)
Harold van Es (Soil and Crop Sciences Section; *soil health/management*)
Robert Schindelbeck (Soil and Crop Sciences Section; *soil health/management*)
Elson Shields (Department of Entomology; *insect, slug, and nematode management*)
Margaret Smith (Plant Breeding and Genetics Section; *grain corn hybrids/small grain variety selection*)
Jaime Cummings (New York State Integrated Pest Management Program; *integrated pest management*)

Abbreviations and Symbols Used in This Publication

A acre	ECemulsifiable concentrate	SPsoluble powder
AI active ingredient	Fflowable	ULVultra-low volume
D dust	Ggranular	Wwetable
DF dry flowable	L liquid	WDGwater-dispersible granules
DG dispersible granule	Ppellets	WPwetable powder
E emulsion, emulsifiable	S soluble	

* Restricted-use pesticide; may be purchased and used only by certified applicators

† Not for use in Nassau and Suffolk Counties

Every effort has been made to provide correct, complete, and up-to-date pest management information for New York State at the time this publication was released for printing (November, 2018). Changes in pesticide registrations, regulations, and guidelines occurring after publication are available in county Cornell Cooperative Extension offices or from the Pesticide Management Education Program web site (pmep.cce.cornell.edu).

Trade names used in this publication are for convenience only. No endorsement of products is intended, nor is criticism of unnamed products implied.

These guidelines are not a substitute for pesticide labeling. Always read and understand the product label before using any pesticide.

The guidelines in this bulletin reflect the current (and past) authors' best effort to interpret a complex body of scientific research, and to translate this into practical management options. Following the guidance provided in this bulletin does not assure compliance with any applicable law, rule, regulation or standard, or the achievement of particular discharge levels from agricultural land.

Cover photo: Winter wheat is grown on 85,000 to 120,000 acres in NY annually with grain yields that average between 63-74bu/acre (NASS 2012-2016). Soft red wheat varieties are most common due to their resistance to sprouting. The photo shows big square bales of wheat straw, baled after wheat harvest at the end of July, in southern Oneida County. The most common uses of wheat straw are as an additive in total mixed rations, or as bedding in the barn. (Photo by: Jeff Miller, Senior Agronomy Educator, CCE Oneida County, NY)

1 PESTICIDE INFORMATION.....	1
1.1 Pesticide Classification and Certification.....	1
1.2 Use Pesticides Safely	1
1.2.1 Plan Ahead	1
1.2.2 Move Pesticides Safely	1
1.2.3 Personal Protective Equipment and Engineering Controls.....	1
1.2.4 Avoid Drift, Runoff, and Spills.....	1
1.2.5 Avoid Equipment Accidents	2
1.2.6 Pesticide Storage	2
1.3 Pollinator Protection.....	2
1.4 New York State Pesticide Use Restrictions.....	2
1.4.1 Restricted-Use Pesticides	2
1.4.2 Additional Use Restrictions	3
1.5 Verifying Pesticide Registration and Restricted-Use Status	3
1.6 Check Label for Site and Pest	3
1.7 Pesticide Recordkeeping/Reporting	3
1.7.1 New York State Requirements.....	3
1.7.2 Federal Private Applicator Recordkeeping Requirements	4
1.8 EPA Worker Protection Standard (WPS) for Agricultural Pesticides.....	4
1.9 Reduced-risk Pesticides, Minimum-risk Pesticides, and Biopesticides	4
1.9.1 Reduced-risk Pesticides	4
1.9.2 Minimum-risk Pesticides	4
1.9.3 Biopesticides.....	5
1.10 FIFRA 2(ee) Recommendations.....	5
2 GENERAL INFORMATION FOR CROP PRODUCTION.....	6
2.1 Introduction.....	6
2.2 New York State Climate	6
2.2.1 Growing Degree Days for Corn and Soybeans	6
2.3 Soil Surveys	7
2.4 Soils and Soil Management Groups of New York	9
2.4.1 Soil Management Group I.....	9
2.4.2 Soil Management Group II.....	10
2.4.3 Soil Management Group III	11
2.4.4 Soil Management Group IV	12
2.4.5 Soil Management Group V	12
2.4.6 Other Areas	13
2.5 Soil Health.....	13
2.5.1 The Cornell Soil Health Test.....	14
2.6 Runoff and Soil Erosion.....	17
2.6.1 Tillage Erosion.....	17
2.6.2 Solving Erosion Problems.....	17
2.6.3 Wind Erosion	18
2.7 Soil Compaction.....	18
2.7.1 Effects of Soil Compaction on Crops.....	20
2.7.2 Diagnosing Plow Layer and Subsoil Compaction.....	20
2.7.3 Addressing Soil Compaction Problems.....	21
2.8 Tillage	23
2.8.1 Full-Width Primary Tillage Systems.....	23
2.8.2 Secondary Tillage	24
2.8.3 Restricted-Tillage Systems.....	24
2.8.4 Deep Tillage	25
2.8.5 Frost Tillage	25
2.8.6 Selecting a Tillage System	26
2.9 Nutrients and the Environment.....	29
2.9.1 Nitrate Leaching Index.....	29
2.9.2 NY Phosphorus Index	30
2.9.3 Manure Spreading Guidelines During Adverse Weather Conditions.....	30

2 GENERAL INFORMATION FOR CROP PRODUCTION *(continued)*

2.10 Soil Testing	34
2.10.1 Why Test Soil?	34
2.10.2 What is a Soil Testing Program?	34
2.10.3 How Does a Soil Test Work?.....	34
2.10.4 How to Collect and Submit a Soil Sample.....	35
2.10.5 How Often Should Samples be Taken?	36
2.10.6 Variable Rate Management.....	36
2.10.7 Interpretation of Soil Test Results	36
2.10.8 Return of Guidelines	37
2.11 Fertilizers	37
2.11.1 Using the Fertilizer Tables.....	37
2.11.2 Fertilizer Materials.....	38
2.11.3 Fertilizer Injury	38
2.11.4 Secondary Macronutrients and Micronutrients.....	39
2.12 Lime Guidelines.....	40
2.12.1 Lime Rate Table.....	41
2.12.2 Lime Applications.....	42
2.12.3 Liming for No-Tillage Production	42
2.12.4 Points to Remember When Using Lime Guidelines Without Buffer pH Data.....	42
2.12.5 Calcium and Magnesium Content of Limestones	43
2.13 Land Application of Sewage Sludges	44
2.13.1 Application Rates.....	45
2.13.2 Nutrients	46
2.13.3 Metals	46
2.13.4 Synthetic Organic Chemicals.....	47
2.13.5 Pathogens.....	47
2.13.6 Sludge Guidelines.....	48
2.14 Integrated Pest Management for Field Crops.....	49
2.14.1 Identification	50
2.14.2 Sampling	50
2.14.3 Analysis	50
2.14.4 Management Alternatives	50
2.14.5 Implementation	50
2.14.6 Evaluation	50
2.15 Integrated Management of Field Crop Diseases	51

3 CORN GUIDELINES.....52

3.1 Planting Techniques.....	52
3.2 Corn Grain Hybrid Selection	53
3.3 Fertilizers for Corn.....	55
3.3.1 Band Rates	56
3.3.2 Efficient Nitrogen Use	56
3.3.3 Nitrogen Status of the Corn Crop	57
3.3.4 Adaptive Management Process.....	60
3.4 Managing Field Corn Pest Problems in New York.....	60
3.5 Managing Diseases of Corn	62
3.5.1 Seed Treatment Fungicides.....	62
3.5.2 Selection of Disease-Resistant Varieties.....	62
3.5.3 Sound Crop Management	62
3.5.4 Foliar Fungicide Application	63
3.6 Managing Insects, Slugs, and Nematodes in Corn.....	64
3.6.1 European Corn Borer.....	65
3.6.2 Northern and Western Corn Rootworms	65
3.6.3 Seed Corn Maggot	66
3.6.4 Cutworms.....	67
3.6.5 Armyworms	67
3.6.6 Wireworms and White Grubs	67

3 CORN GUIDELINES *(continued)*

3.6 Managing Insects, Slugs, and Nematodes in Corn <i>(continued)</i>	
3.6.7 Potato Stem Borer and Hopvine Borer.....	68
3.6.8 Western Bean Cutworm	68
3.6.9 Slugs.....	68
3.6.10 Nematodes.....	68
3.7 Managing Weeds in Corn.....	70
3.7.1 Weed Management Methods.....	70
3.7.2 Factors Affecting Soil-Applied Herbicides	70
3.7.3 Herbicide Resistance Management	71
3.7.4 Weed Control in Zone/No-Tillage Corn	71
3.7.5 Prepackaged Herbicide Mixtures	72
3.7.6 *Atrazine Application Rates	72

4 FORAGE CROPS GUIDELINES **83**

4.1 Forage Varieties	83
4.1.1 Alfalfa	83
4.1.2 Birdsfoot Trefoil	83
4.1.3 Red Clover	84
4.1.4 Alsike Clover	84
4.1.5 Crownvetch.....	84
4.1.6 Results from Forage Trials harvested in 2017 (trials sown in 2014, 2015, and 2016)	84
4.2 Perennial Grasses	89
4.2.1 Reed Canarygrass.....	89
4.2.2 Timothy.....	89
4.2.3 Smooth Brome grass	89
4.2.4 Orchardgrass	89
4.2.5 Ryegrasses.....	89
4.2.6 Tall Fescue	90
4.2.7 Festulolium	90
4.2.8 Perennial Forage Variety Selection.....	90
4.3 Annual Crops for Forage.....	95
4.3.1 Sudangrass and Forage Sorghums.....	95
4.3.2 BMR Sorghum-Sudangrass Hybrids.....	95
4.3.3 Anti-Quality Components	96
4.3.4 Millets	96
4.3.5 Oats and Spring Triticale.....	96
4.3.6 Winter Wheat, Barley, Rye, and Triticale	96
4.4 Planting Perennial Legumes and Grasses.....	97
4.4.1 Seedbed Preparation.....	97
4.4.2 Date of Seeding	97
4.4.3 Inoculation	97
4.4.4 Fertilizers and Lime for Legume Establishment	97
4.4.5 Companion Crops	97
4.4.6 Species Selection.....	97
4.5 Harvest Schedules	97
4.5.1 Alfalfa	97
4.5.2 Perennial Grasses	98
4.6 Forage Fertilization	98
4.6.1 Considerations for Manure Use for Alfalfa-Grass production	99
4.7 Forages for Conservation	101
4.8 Managing Alfalfa Pest Problems in New York.....	102
4.9 Managing Diseases of Perennial Forage Legumes.....	103
4.9.1 Vascular Wilts.....	104
4.9.2 Root, Crown, and Stem Rots.....	104
4.9.3 Leaf and Stem Blights	104
4.9.4 Seed Fungicides	104
4.9.5 Selection of Disease-Resistant Varieties	104

4 FORAGE CROPS GUIDELINES *(continued)*

4.9 Managing Diseases of Perennial Forage Legumes <i>(continued)</i>	
4.9.6 Sound Stand Management	105
4.10 Managing Insects in Forage Crops.....	105
4.10.1 Alfalfa Weevil	105
4.10.2 Potato Leafhopper.....	106
4.10.3 Alfalfa Snout Beetle.....	108
4.11 Weed Control in Forages	110
4.11.1 Herbicide Resistance Management.....	110
4.12 Minimum-Tillage Forage Seedings.....	114
4.12.1 Vegetation Control.....	114
4.12.2 Seeding Methods.....	114
4.12.3 Insect Control.....	114
4.13 Perennial Forages for Pasture	114
4.13.1 Ladino White Clover (<i>Trifolium repens</i> L.).....	115
4.13.2 Kentucky Bluegrass (<i>Poa pratensis</i> L.)	115
4.13.3 Tall Fescue (<i>Festuca arundinacea</i> Schreb).....	115
4.13.4 Use of Pastures	115
4.13.5 Grass-Legume Mixtures.....	115
4.14 Weed Control in Grass Pastures.....	116
4.14.1 Herbicide Resistance Management.....	116

5 SMALL GRAIN CROPS GUIDELINES.....118

5.1 Planting Techniques.....	118
5.1.1 Winter Grains.....	118
5.1.2 Spring Grains	118
5.2 Variety Selection.....	118
5.2.1 Winter Wheat.....	118
5.2.2 Hybrid Winter Rye.....	119
5.2.3 Spring Oat.....	119
5.2.4 Winter Malting Barley	119
5.2.5 Spring Malting Barley	120
5.2.6 Triticale.....	120
5.2.7 Spelt.....	120
5.2.8 Buckwheat	120
5.2.9 Other Grain-Type Crops	120
5.3 Why We Recommend Certified Seed	124
5.4 Growth Stages in Small Grains.....	124
5.4.1 Winter Wheat.....	124
5.4.2 Spring Grains	125
5.5 Fertilizers for Small Grains.....	125
5.5.1 Lodging Control in Wheat or Barley	126
5.6 Managing Winter Wheat Pest Problems in New York.....	127
5.7 Managing Diseases of Small Grain Cereals.....	128
5.7.1 Selection of Disease-Resistant or Disease-Tolerant Varieties	129
5.7.2 Seed Treatment	129
5.7.3 Foliar Fungicides	129
5.7.4 Guidelines for Wheat Fungicide Decisions	129
5.7.5 Sound Agronomic Practices.....	130
5.8 Managing Insects in Small Grains	132
5.8.1 Armyworms	132
5.8.2 Cereal Leaf Beetle	133
5.8.3 Hessian Fly	133
5.9 Weed Control in Small Grains	134
5.9.1 Herbicide Resistance Management.....	134

6 SOYBEAN GUIDELINES.....	136
6.1 Soybean Maturity Group Selection	136
6.2 Soybean Planting.....	136
6.3 Managing the Crop.....	136
6.4 Managing Soybean Pest Problems in New York.....	137
6.5 Managing Diseases of Soybeans	140
6.5.1 Fungicidal Seed Treatment	140
6.5.2 Foliar Fungicides.....	140
6.5.3 Asian Soybean Rust	140
6.5.4 Soybean Cyst Nematode	140
6.6 Managing Insects in Soybeans	142
6.6.1 Seed Corn Maggot	142
6.6.2 Soybean Aphid.....	143
6.6.3 Two-spotted Spider Mites	143
6.7 Weed Control in Soybeans	144
6.7.1 Herbicide Resistance Management	144
7 TOTAL VEGETATION CONTROL.....	150
7.1 Total Vegetation Control.....	150
8 APPENDIX	151
8.1 Trade and Common Names of Field Crop Pesticides.....	151
8.2 Herbicide Site of Action Groups and Mode/Site of Action Information.....	154

1 Pesticide Information

1.1 Pesticide Classification and Certification

The Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) created two classifications of pesticides – general-use and restricted-use. **General-use pesticides** may be purchased and used by anyone. **Restricted-use pesticides can** only be purchased by a certified applicator. Restricted-use pesticides must also be used by a certified applicator or someone under their supervision.

The same federal law that classifies pesticides divided applicators into two groups: private and commercial.

Private applicators use or supervise the use of pesticides to produce agricultural commodities or forest crops on land owned or rented by the private applicator or their employer. A farmer must be certified as a private applicator in order to purchase and use restricted-use pesticides on agricultural commodities. (No certification is needed if a farmer does not use restricted-use pesticides.)

A **commercial applicator** uses or supervises the use of pesticides for any purpose or on any property not covered by the private applicator classification. In New York, a commercial applicator must be certified to purchase or use any pesticide whether it is general- or restricted-use.

Information about pesticide certification and classification is available from your Cornell Cooperative Extension office (cce.cornell.edu/localoffices), regional NYSDEC pesticide specialist (www.dec.ny.gov/about/558.html), the Pesticide Applicator Training Manuals (<https://www.cornellstore.com/books/cornell-cooperative-ext-pmep-manuals>), or the Pesticide Management Education Program (PMEP) at Cornell University (psep.cce.cornell.edu).

1.2 Use Pesticides Safely

Using pesticides imparts a great responsibility on the user to protect their health and that of others and to protect the environment. Keep in mind there is more to “pesticide use” than the application. Pesticide use includes mixing, loading, transporting, storing, or handling pesticides after the manufacturer’s seal is broken; cleaning pesticide application equipment; and preparing a container for disposal. These activities require thoughtful planning and preparation. They are also regulated by state and federal laws and regulations intended to protect the user, the community, and the environment from any adverse effects pesticides may cause.

1.2.1 Plan Ahead

Many safety precautions should be taken *before* you actually begin using pesticides. Too many pesticide applicators are dangerously and needlessly exposed to pesticides while they are preparing to apply them. Most

pesticide accidents can be prevented with informed and careful practices. **Always read the label on the pesticide container before you begin to use the pesticide.** Make sure you understand and can follow all directions and precautions on the label. Be prepared to handle an emergency exposure or spill. Know the first aid procedures for the pesticides you use.

1.2.2 Move Pesticides Safely

Carelessness in transporting pesticides can result in broken containers, spills, and contamination of people and the environment. Once pesticides are in your possession, you are responsible for safely transporting them. Accidents can occur even when transporting materials a short distance. You are responsible for a pesticide accident so take every effort to transport pesticides safely. Be prepared for an emergency.

1.2.3 Personal Protective Equipment and Engineering Controls

Personal protective equipment needs depend on the pesticide being handled. ***Required personal protective equipment (PPE) are listed on pesticide labels.*** The required PPE are based on the pesticide’s toxicity, route(s) of exposure, and formulation. Label required PPE are the minimum that must be worn during the pesticide’s use. Pesticide users can always wear more protection than the label requires.

The type of protective equipment used depends on the type and duration of the activity, where pesticides are being used, and exposure of the handler. Mixing/loading procedures often require extra precautions. Studies show you are at a greater risk of accidental poisoning when handling pesticide concentrates. Pouring pesticide concentrates from one container to another is the most hazardous activity.

Engineering controls are devices that help prevent accidents and reduce a pesticide user’s exposure. One example is a closed mixing/loading system that reduces the risk of exposure when dispensing concentrated pesticides. Consult the product label for more information on using engineering controls in place of PPE.

1.2.4 Avoid Drift, Runoff, and Spills

Pesticides that move out of the target area can injure people, damage crops, and harm the environment. Choose weather conditions, pesticides, application equipment, pressure, droplet size, formulations, and adjuvants that minimize drift and runoff hazards. See product labels for specific application and equipment requirements.

2 General Information for Crop Production

2.1 Introduction

This publication includes the most up-to-date information on growing field crops in New York, drawn from Cornell research, extension demonstrations, and on-farm experience. It has been designed as a practical guide for farmers, for merchants who provide sales and services to producers, and for others who advise them. Our aim is to supply the best information available to help those who make management decisions. We do not consider this a cookbook but rather a source of practical information to use in the development of sound planning and good management.

In any statewide publication, we must deal with a spectrum of crop environments; information and guidelines must cover general farm situations. Though we have tried to make these as specific as possible for various conditions in New York, each farmer must determine how these varieties and practices will work on his or her farm. The information in this publication should be considered general rules. Additional information is available in the publication **Cornell Field Crops and Soils Handbook**, revised in 1987 and available through Cornell Cooperative Extension offices or directly from the Section of Soil and Crop Sciences at Cornell, 607-255-2177. For further information on any topic in this booklet, you may contact your local Cornell Cooperative Extension office or write to the Section of Soil and Crop Sciences Extension Office, 237 Emerson Hall, Cornell University, Ithaca, NY 14853.

2.2 New York State Climate

2.2.1 Growing Degree Days for Corn and Soybeans

Crop plants require heat from their atmospheric environment to develop, grow, and mature. The effect of this heat is cumulative as the growing plant progresses through its life cycle.

Temperature is an indirect measure of the heat available in the atmosphere. Heat sufficient to cause growth and development in a plant is indicated when the daily mean temperature warms to a certain level, called the base or threshold temperature. Below (cooler than) this level there is essentially no growth. Different species of crop plants have different base temperatures. Corn and soybeans have a base temperature of 50°F.

The growing degree day—sometimes called a heat unit—has become a useful indirect measure of the heat available for growth and development of corn and soybeans. In the 86/50 method it is assumed that for corn and soybeans, growth increases linearly from 50°F to 86°F, at which peak growth occurs, and growth remains at peak for temperatures above 86°F. The maximum temperature for the day is set at

an upper limit of 86°F, and the minimum temperature is set at the lower limit of 50°F. On each day of the growing season the crop receives a number of growing degree days equal to the number of degrees that the daily adjusted mean temperature is higher (warmer) than the 50°F base temperature. Growing degree days are then accumulated each day as the crop progresses toward maturity.

To calculate the daily growing degree days for your farm, first, determine the adjusted mean air temperature for each 24-hour day during the growing season. For a day with a high temperature of 60°F and a low of 40°F, for example, the low temperature would be set at the lower limit of 50°F. The adjusted mean temperature for the day would be 55°F. Subtracting 50°F, the base temperature for corn and soybeans, from the mean temperature gives 5 growing degree days for that day. If, on the other hand, the high temperature for a given day is 90°F and the low is 66°F, the high temperature would be set at the upper limit of 86°F. The adjusted mean temperature for the day would be 76°F. Subtracting 50°F, the base temperature for corn and soybeans, from the mean temperature gives 26 growing degree days for that day. On any day that the adjusted mean temperature is 50°F or colder, the number of growing degree days is recorded as zero.

Records are kept for each day of the growing season, from the first frost-free day in the spring through the last frost-free day in the fall. By adding together the growing degree days supplied each day, the accumulated total for the frost-free growing season is determined.

The distribution of average accumulative growing degree days in New York State is presented in Figure 2.2.1. These data, applying to the freeze-free season, were determined from temperature records kept by numerous weather stations around the state during a 30-year period ending in 1980.

2.2.1.1 How to Use Growing Degree Days

Early corn hybrids and short-season soybean varieties need fewer growing degree days than late corn hybrids and long-season soybean varieties to grow and mature.

Use the map (Figure 2.2.1) to determine the growing degree days available for corn and soybean growth in your locality. You can then choose corn hybrids and soybean varieties suited to your vicinity from the groups listed in sections 3.2: Corn-Hybrid Selection and 6.1: Soybean Varieties. You may need to make adjustments to fit local differences in elevation or frost susceptibility.

3 Corn Guidelines

High-yielding corn requires moderately well-drained or well-drained soil with a pH above 6.0 as well as timely and skillful management practices. Management practices to consider carefully include planting techniques, hybrid selection, fertilization, and control of insects, weeds, and diseases. Correct management of all these practices is essential for maximum economic yield.

3.1 Planting Techniques

Early planting usually, but not always, results in maximum corn yields. Under central and western New York conditions, corn planted in late April or early May typically out yields either grain or silage corn planted after mid-May (Figure 3.1.1). Early-planted corn also matures earlier, resulting in lower moisture and grain drying costs at harvest, and lodges less. A general guideline for the best time to begin planting corn is about 10 days before the average date of the last 32°F temperature in the spring. If soil conditions are too wet at this time, wait until soil conditions improve. Corn planted in late May under dry soil conditions will consistently out yield corn planted in late April under wet soil conditions. Conversely, if it is warm and dry any time after April 15th in central/western NY, corn growers should be ready to begin planting. Modern corn hybrids tolerate cold soil conditions and seed treatments protect corn from soil pest problems under extended emergence time due to cold soil temperatures. Planting depths of about 1.5 inches for silty clay or clay loam soils and 1.75 to 2.0 inches for silt loam and gravelly loam soils are recommended for April or early May-planted corn. Planting depths of about 1.75 to 2.0 inches for silty clay or clay loam soils and 2.0 to 2.5 inches for silt loam and gravelly loam soils are recommended for most planting dates in May. If soil conditions are dry in the top 2 inches in late May and early June, corn can be safely planted to a depth of 3 inches on silt loam and gravelly loam soils.

To achieve the full yield potential of an early planting date, full-season hybrids (hybrids that match the growing degree days in a region) are necessary (Figure 3.1.1). After the first or second week of May, however, the yield advantage of full-season vs. medium-season hybrids decreases when planted for grain. Furthermore, full-season hybrids may not mature, resulting in low test weight, and/or will have high grain moisture at harvest, if planted after the second week of May. Therefore, for grain production, full-season hybrids should be planted only in late April or during the first 2 weeks of May. For silage production, full-season hybrids can be planted until about May 20. Growers should not plant more than 30% of their crop to full-season hybrids. The majority of corn acreage (~60%) should be planted to medium-season hybrids (100 and 200 growing degree days less than the growing degree days in a region for silage and grain, respectively). If planting must be delayed until early June, early-season hybrids (300-400 growing degree days

less than the growing degree days in a region for silage and grain, respectively) are recommended.




Figure 3.1.1. Effect of planting date on grain yields.

The optimal corn population depends on soil type, hybrid selection, and crop use. For many New York soils (well-to moderately well-drained to somewhat poorly drained silt or clay loams), numerous Cornell experiments have shown that modern hybrids still require a harvest population of only 26,000 to 28,000 plants per acre for maximum economic grain yields (Table 3.1.1). Droughty soils, however, cannot support high populations, and plant populations should be adjusted downward (Table 3.1.1). Likewise, hybrids differ in their response to plant populations, so hybrid selection should influence whether the harvest population is at the high or low end of the recommended range for each particular soil condition (Table 3.1.1). Also, most hybrids require higher harvest populations for silage than for grain production, about 5,000 more plants per acre (Table 3.1.1).

Planting date, tillage practices, pest problems, planter performance, and hybrid selection influence actual corn populations obtained in the field. To compensate for potential problems, it should be assumed that only 90 percent of the kernels planted actually emerge and survive to become harvestable plants in the fall. To obtain 27,000 plants per acre at harvest on a moderately well-drained silt-loam soil, the planting rate should be about 30,000 plants per acre (27,000 divided by 0.90). In some situations such as a no-till situation or an April planting date, it should be assumed that only 85 percent of the kernels will emerge and survive. The planting rate in these situations on a moderately well-drained silt-loam soil should be about 31,765 plants per acre (27,000 divided by 0.85).

Table 3.4.1. Field corn pests and crop monitoring activities.

Routine	Occasional
Early Season (emergence to sixth leaf stage)	
Population counts (establishment problems, seed and seedling pests), weeds, cutworm, seedling diseases, damping off; seed corn maggot, watch for “occasional” pests	White grubs, common armyworm, slugs, common stalk borer, billbug, stink bug, corn flea beetle, foliar diseases, nutrient deficiencies, environmental stress (e.g., cold weather, excessive moisture or drought), birds, deer, hail)
Mid-Season (sixth leaf stage through early silk stage)	
Presidedress nitrogen sampling; watch for “occasional” pests	Weed escapes, common armyworm, European corn borer, western bean cutworm, foliar blights, anthracnose, common stalk borer, potato stem borer, hopvine borer, vertebrate pests, nutrient deficiencies, environmental stress, hail
Pollination (throughout silking and early grain fill stages)	
Corn rootworm, northern corn leaf blight, northern corn leaf spot, eyespot, common rust; watch for “occasional” pests	European corn borer, western bean cutworm, foliar blights, stalk rots, vertebrate pests, nutrient deficiencies, environmental stress, hail
Late Season (harvest and postharvest)	
Ear mold assessment, yield checks, weed inventory survey, soil sampling	European corn borer, foliar blights, stalk rots, ear molds, vertebrate pests, environmental stresses, lodging assessment


Figure 3.4.1. Field Corn IPM Scouting Calendar.

Table 3.7.3. Chemical weed control in corn.

Weed Situation	Amount of Product(s) per Acre	Remarks and Limitations
Conventional corn hybrids (continued)		
PRE Programs Annual grass and broadleaf weeds (continued)	3.0-3.5 qt. *†Lexar EZ (continued)	organic matter is 3% or greater. *†Lexar EZ provides twice as much *atrazine/A as *†Lumax EZ (see Table 3.7.6 for the amount of *atrazine at labeled rates of each product). This additional *atrazine should improve common ragweed control. *†Lexar EZ is not for use in Nassau and Suffolk Counties.
	3 pt. Prowl H2O or 3.6 pt. Prowl 3.3 EC + 1 qt. *†Atrazine 4L ¹	GROUP 3 and 5 HERBICIDES • Apply preemergence. Prowl should not be applied preplant incorporated for corn. Plant corn at least 1 1/2 in. deep in fields with adequate seedbed preparation to provide good coverage of the corn seed. Good choice if velvetleaf or triazine-resistant lambsquarters are problems.
POST Programs Emergent annual grass and broadleaf weeds	1.5 oz. Steadfast Q + 4 fl. oz. Banvel or 4 fl. oz. Clarity or 4 fl. oz. DiFlexx	GROUP 2 and 4 HERBICIDES • Apply early postemergence with crop oil concentrate or nonionic surfactant and ammonium nitrogen fertilizer to hybrids with a relative maturity (RM) rating of 77 days or more. For best results, apply when annual weeds are 1-2 in. tall and before corn is 12 in. tall. A good choice if foxtails or fall panicum and annual broadleaf weeds have emerged. Not very effective against crabgrass. Rotational interval is 10 months or longer (depending on soil pH) for alfalfa, clovers, and several other crops. Do not apply Steadfast Q if corn was treated with Counter 15G (any application method).
	0.75 fl. oz. Impact + 1 pt *†Atrazine 4L	GROUP 27 and 5 HERBICIDES • Apply early postemergence with methylated seed oil (MSO) or crop oil concentrate and ammonium nitrogen fertilizer when annual weeds are 1 - 2 in. tall. May be tank mixed with half rates of preemergence herbicides shown above under PRE Programs for added residual weed control. Use COC if tank mixing with Prowl rather than MSO. Rotation interval following Impact application is 3 months for small grains and 9 months for alfalfa, peas, potatoes, and soybeans.
	3 fl. oz. Laudis + 1 pt *†Atrazine 4L	GROUP 27 and 5 HERBICIDES • Apply early postemergence with methylated seed oil (MSO) and ammonium nitrogen fertilizer when annual weeds are 1 - 3 in. tall. May be tank mixed with half rates of preemergence herbicides shown above under PRE Programs , except Prowl combinations, for added residual weed control. Rotation interval following Laudis application is 4 months for small grains, 8 months for soybeans, and 10 months for alfalfa, potatoes, and snap beans.
	3 fl. oz. *†Capreno + 1 pt *†Atrazine 4L	GROUP 2, 5, and 27 HERBICIDES • Apply early postemergence with crop oil concentrate and ammonium nitrogen fertilizer when annual weeds are 1 – 3 in. tall. Rotational interval is 4 months for wheat, 10 months for barley, soybeans and sweet corn, and 18 months for alfalfa, oats, and some other crops.
Problem Weed Quackgrass	22–44 fl. oz. Roundup PowerMax or 24–48 fl. oz. Durango DMA or 24–48 fl. oz. Touchdown Total	GROUP 9 HERBICIDES • Apply these or other glyphosate products as a foliar spray in the spring or fall when quackgrass is at least 8 in. tall (4- to 5-leaf stage and actively growing). Fall applications should not be made in fields that have been tilled during the summer or mowed after August 15. Delay tillage for at least 3 days after spraying. The low rates may not provide long-term quackgrass control.
	1.5 oz. Steadfast Q or 0.76 oz. *Beacon	GROUP 2 HERBICIDES • Apply postemergence when quackgrass is 4 to 8 in. tall and corn is between 4 and 20 in. A nonionic surfactant or a crop oil concentrate must be used. Do not apply Steadfast Q or *Beacon if corn was treated with Counter 15G (any application method). Rotational intervals for alfalfa and several other crops are 10 months or longer (depending on soil pH) for Steadfast Q and are 8 months for *Beacon.

4 Forage Crops Guidelines

4.1 Forage Varieties

4.1.1 Alfalfa

Often called the queen of the forages, alfalfa tops all other perennial forage crops as a producer of homegrown feed. High-yielding and versatile, alfalfa serves well for hay, silage, green chop, and pasture. It produces high-protein and palatable feed, which livestock like and do well on. Alfalfa also fills an important role in crop rotations, improving soil structure and building soil fertility for future grass and grain crops.

Alfalfa is a deep-rooted, drought-tolerant crop that does best on deep, well-drained soils. Alfalfa also needs a well-limed soil; it gives top performance on soils with pH levels of 6.5 or higher. It does poorly on acidic soils, and soil acidity is often noted as the major limiting factor on alfalfa growth in New York. Acidic soils must be limed to a pH of 6.5 or higher to maintain high-yielding alfalfa stands.

On well-drained soils, alfalfa can produce high yields for many years, but it will yield poorly and die soon on poorly drained soils. Tile and other drainage aids can improve the soil's ability to grow good alfalfa. Trefoil and red clover offer better choices for good production on sites with poor or spotty drainage patterns.

Alfalfa seedlings need phosphorus and potassium at planting time. Older stands need topdressing to maintain high yields. An ample fertility program provides nutrients for recovery after harvest, good winter survival, and high yields. Phosphorus and potassium are musts, but nitrogen rarely, if ever, pays on alfalfa because nitrogen-fixing bacteria in root nodules can provide enough nitrogen for top yields. For details on fertilizer suggestions, see Table 4.6.1.

Insect pests cause sporadic damage in alfalfa, varying with season and locality. Potato leafhopper feeding can lower second-cut yields in some years. The alfalfa weevil and blotch leaf miner, formerly serious, are now largely controlled through introduced insect parasites and predators. The alfalfa snout beetle can cause severe damage in the several counties where it occurs. Check control guidelines in the section “Management of Insects in Forage Crops (section 4.10).”

New York alfalfa trials test yield of new varieties (Table 4.1.1 to 4.1.6). Modern alfalfa varieties have been bred for resistance to five or more diseases that can thin alfalfa stands in New York. These diseases include **bacterial wilt**, caused by bacteria present in most New York alfalfa soils; **Verticillium wilt**, a soilborne disease that can kill susceptible plants in their second or third year; **Phytophthora** root rot, caused by a soilborne water mold often found in wet areas of fields; **anthracnose**, found in warmer areas of the state, particularly the Hudson Valley; and **Fusarium wilt**, common in New York soils and may occur but is not documented as a

widespread problem in New York. *Phytophthora* hits hardest in the seedling year, and the other diseases affect mature stands in their second and third years of production.

Check for variety reactions to these specific diseases as well as for yield and fall dormancy ratings. Choose varieties that are listed as R (resistant) or HR (highly resistant) for diseases found in your area. View resources online and table 4.1.6 to determine disease resistance ratings and other characteristics for alfalfa varieties. View tables 4.1.1 to 4.1.3 for fall dormancy ratings and relative yield comparisons. Both *Aphanomyces* root rot and pea aphid occur here but the value of varietal resistance may not be established for these and some other pests.

Several varieties have been developed at Cornell for specific adaptation to New York State conditions. These include ReGen, Ezra, N-R-Gee, Seedway 9558 SBR (selected for resistance to alfalfa snout beetle) and SW315LH (selected for resistance to potato leafhoppers).

Improved feeding value has been a goal of alfalfa breeders for years. Several recent varieties have been released with claims of improved feeding quality. Our tests show that minor differences in feeding quality do exist. However, effects on milk production have yet to be established. Timely cutting and leaf-saving harvest practices are far more important in affecting forage quality than leaf or plant type. Choose varieties with strong disease resistance and high yield potential that are well adapted to your farm and needs. Optimal yield and forage quality is at the one-tenth bloom stage.

New leafhopper-resistant varieties are available that have improved resistance and agronomic characteristics (see Table 4.1.2). Leafhopper resistance comes from fine hairs on stems and leaves, and results in significantly lower numbers of hoppers in resistant alfalfa stands compared to conventional alfalfa. Resistant varieties will surpass other strains when leafhopper pressure is heavy. Spraying in the seeding year may still pay under heavy hopper pressure.

4.1.2 Birdsfoot Trefoil

Birdsfoot trefoil is a long-lived legume with high yield potential on slightly acidic soils with drainage less than the best for alfalfa. Trefoil also does well as perennial forage on hard-to-plow meadows and pastures. Trefoil is bloat free, and no case of bloat has ever been recorded in animals grazing on trefoil. On fields where drainage is a problem, trefoil can out yield alfalfa and outlive red clover by many years. Birdsfoot trefoil should be planted with a perennial forage grass and at harvest time, leave 5 to 6 inches of stubble to allow for regrowth of the trefoil.

PARDEE birdsfoot trefoil is a vigorous, upright, hay-type variety. It resists the *Fusarium* wilt disease that often kills

Table 4.1.4. 2017 Yields of Red Clover Varieties in trials planted in 2014, 2015, and 2016 in Central New York (CNY – Ithaca). Company contact information in Table 4.2.4.

T/A = tons per acre dry matter

Red Clover	Company	Third Prod Yr. Yield	Second Prod. Yr. Yield	First Prod. Yr. Yield
EVOLVE	DLF Pickseed USA Inc.			99
FF9615	La Crosse Seed		96	
FSG 402	Seedway		103	
Gallant Red	Preferred Seed Co.	106		
GO-MER	Grassland Oregon	89		
Marathon (check)	Public Check	97	99	97
# Trial Entries		9	5	8
Check Variety Mean (T/A/Yr)		3.61	5.10	6.89
Trial Mean (T/A/Yr)		3.53	4.91	6.76

Table 4.1.5. Alfalfa trials harvested for yield in 2017: trial location, NY region, Seeding year, soil series, elevation, number of harvests and number of production years.

Trial Location, NY Region*, Seeding Year	Soil series, elevation, # of harvests in 2017, # of production years
Ithaca, CNY, 2014	Rhinebeck silt loam, 970 ft., 3 harvests in third production year
Richfield Springs, ENY, 2014	Honeoye silt loam 1400 ft., 3 harvests in third production year
Ithaca, CNY, 2015	Williamson silt loam, 1000 ft., 3 harvests in second production year.
Chazy, NNY, 2015	Raynham variant silt loam, 185 ft., 2 harvests in second production year.
Ithaca, CNY, 2016	Erie Chippewa channery silt loam, 1054 ft. 3 harvests in first production year.
Cobleskill, ENY, 2016	Barbour Tioga fine sandy loam, 1170 ft., 3 harvests in first production year.
Ithaca, CNY, 2017	Williamson silt loam, 1000 ft., 0 harvests in seeding year.
Geneva, WNY, 2017	Honeoye and Lima loam, 486 ft., 0 harvests in seeding year.

*CNY – Central New York, ENY – Eastern NY, WNY – Western NY, NNY – Northern New York.

Table 4.1.6: Alfalfa Cultivar FeaturesFor more information log on to the Web:<http://plbrgen.cals.cornell.edu/research-extension/forage-project/ny-forage-yield-results>

Cultivars listed are currently tested in Cornell Alfalfa Trials. Yield data for cultivars in new trial seedings will be available next year.

Alfalfa Cultivar	Marketing Company	FD	Disease Resistance Ratings*					Marketing Co. Contact Information	
			BW	VW	FW	AN	PRR	Phone	Web or E-mail Address
59W205	BrettYoung	5	R	HR	HR	HR	HR	1-800-665-5015	www.brettyoung.ca/
KINGBIRD	Blue River Organic Seed	5	HR	HR	HR	HR	HR	1-800-370-7979	www.blueriverorgseed.com
RED FALCON	Blue River Organic Seed	4	HR	HR	HR	HR	HR		
ROBIN	Blue River Organic Seed	5	HR	HR	HR	HR	HR		
RR APHATRON 2XT	CROPLAN	4	HR	HR	HR	HR	HR	1-651-765-5710	www.croplan.com
REBOUND 6XT	CROPLAN	4	HR	HR	HR	HR	HR		
HYBRIFORCE-3400	Dairyland Seed Co.	4	HR	HR	HR	HR	HR	1-800-236-0163	www.dairylandseed.com
HYBRIFORCE-3430	Dairyland Seed Co.	4	HR	HR	HR	HR	HR		

Table continues on next page.

5 Small Grain Crops Guidelines

Small grains, which include winter and spring wheat, winter and spring barley, oats, and rye, play an important role in crop rotations on many New York farms. Under good soil conditions and management practices, small grains can produce profitable yields of grain for the cash market or farm feeding. Equally important is the value of the straw crop.

Oats and rye tolerate acid or poorly drained soils better than wheat or barley does. Nevertheless, maximum yields of both crops are attained on moderately well-drained or well-drained soils with a pH above 5.8. For maximum wheat production, wheat must be cropped on moderately well-drained or well-drained soils with a pH above 6.0. Barley requires well-drained soils with a pH above 6.3, the same as needed for alfalfa production.

5.1 Planting Techniques

5.1.1 Winter Grains

Winter wheat should be planted with a grain drill to a depth of 1 to 1-1/2 inches during the couple of weeks after the Hessian fly-free date. The optimal planting is thus from mid-September until early October in most regions of winter wheat production. Depending upon the fall or winter conditions, wheat can be successfully planted until early November but at a lower yield potential. Soft white winter wheat has a broad optimum seeding rate range and rates of about 120 pounds or 2 bushels per acre usually result in the highest grain and straw yields. If planting is delayed beyond early October, the optimal rate is 150 pounds or 2-1/2 bushels per acre. Soft red winter wheat also has a broad optimum seeding rate range and rates between 1,000,000 and 1,300,000 seeds per acre result in highest grain yields when planted in September and about 1,500,000 seeds/acre for highest straw yields. If planting is delayed after mid-October, soft red winter wheat should be seeded at rates of 1,500,000 seeds for acre if just for grain and 1,800,000 seeds/acre if the straw is also harvested.

Barley is less hardy than wheat and is not susceptible to Hessian fly. Winter barley can thus be planted a few days earlier than wheat, that is, from September 10 to September 20. Because barley is very susceptible to barley yellow dwarf virus, planting before this time is strongly discouraged. It is best to sow the seed with a grain drill at a depth of 1 to 1-1/2 inches. Seeding rates should be in the 96 to 120 pounds per acre or 2- to 2-1/2-bushel range.

Rye is the hardiest of all winter grains and thus can be successfully established with an early to mid-October planting date. For seed production, rye should be sown with a grain drill at a depth of 1 to 1-1/2 inches. The seeding rate should be in the 110 pounds or 2-bushel range.

5.1.2 Spring Grains

Spring grains should be sown as early in the spring as possible. In central New York, a yield decrease of about 1 bushel per acre per day can be expected in oats and barley for each day the crop is planted after April 15. With spring wheat, a yield loss of about 1/2 bushel per acre per day can be expected if planting occurs after April 15. All spring grains should be sown with a grain drill to a depth of 1 to 1-1/2 inches. The optimal seeding rate for oats is 96 pounds or 3 bushels per acre, whereas spring barley and spring wheat do best at 2 bushels per acre. If oats or barley is to be used in forage seeding, seeding rates should be reduced by 50 percent.

See the *Cornell Field Crops and Soils Handbook* for more detailed planting information.

5.2 Variety Selection

5.2.1 Winter Wheat

Wheat is an important cash crop in central and western New York. Most New York wheat is classified as soft red winter wheat, but some soft white winter wheat is also grown. Millers use the soft wheats to produce high-quality, low-protein flours for use in pastries, crackers, cookies, and breakfast cereals. Soft red wheats are inherently more resistant to pre-harvest sprouting than soft white wheats.












Winter wheat varieties are tested every year in Cornell trials, and results of multiyear evaluations are shown for soft white wheat varieties from Cornell's breeding program in Table 5.2.1 and for both Cornell and commercial soft red wheat varieties in Table 5.2.2. Please note the following points when using these tables. Varieties are in order from those that have been tested the longest to those most recently entered into the testing program. For each trait, the number of years of data used to assess that trait are noted at the top of the table. The more years of evaluation, the more precise the data will be. **The table includes only varieties that have been tested for at least two years in Cornell trials.** All the winter wheat varieties reported in these tables are good options for New York growers. Their yields are good and all have acceptable milling and baking quality, test weight, and lodging resistance.



5.2.1.1 Soft White Winter Wheat

Only varieties developed by Cornell University's soft white wheat breeding program are evaluated in Cornell trials at this time. Results of variety evaluations are reported in Table 5.2.1. Special traits of some of these varieties are noted below, but recall that **all the varieties listed in the Table are good options for New York growers.**

CALEDONIA is a good yielder with excellent standability. It has attractive, light-colored straw.

Figure 5.6.1. Winter wheat IPM scouting calendar.

	September	October	November	Dec.-Mar.	April	May	June	July
								
					stand count/assessment			
Weeds					weed survey			
Diseases								
					monitor closely for virus diseases			
								
						monitor closely for fungal diseases		
Insects								
	aphids/Hessian fly					aphids		
								
						cereal leaf beetle		
								
					cutworm			
						armyworm/fall armyworm		
								
						European corn borer, grasshoppers		
	September	October	November	Dec.-Mar.	April	May	June	July

Key:  Preferred  May vary with season

5.7 Managing Diseases of Small Grain Cereals

Diseases are often a major yield constraint in the production of wheat, oats, and barley in New York State. Many disease-induced losses have been accepted by growers because of either a lack of acceptable control practices or a lack of disease detection. Disease management has been accomplished to some extent through the use of sound cultural practices, seed-treatment fungicides, and varieties resistant (or at least less susceptible) to certain diseases. With renewed interest in intensive management, the severity and importance of diseases have increased, and greater emphasis has been placed on the use of newly available foliar fungicides.

Overall, the most important diseases are fungal foliar diseases. These include *Stagonospora nodorum* blotch, *Septoria tritici* blotch, tan spot, powdery mildew, and leaf

rust on wheat; crown rust, *Septoria* blight, and leaf blotch on oats; and powdery mildew, spot blotch, net blotch, scald, and leaf rust on barley. Foliar diseases, if severe during the first few weeks after flowering, can drastically reduce grain yields.

Soilborne fungal diseases affecting the roots, crowns, and lower stems occur on cereals in New York but are seldom severe. This is probably because small grains seldom follow each other in rotational sequence in New York and the same cereal species is almost never continuously cropped. Diseases such as take-all, *Fusarium* root and crown rot, eyespot and other foot rot diseases can reduce cereal yields under certain conditions in New York.

Yellow dwarf, caused by aphid-transmitted viruses collectively called barley yellow dwarf and cereal yellow dwarf viruses, affects all cereals in New York and can be a serious problem. Wheat spindle streak mosaic, caused by a

6 Soybean Guidelines

6.1 Soybean Maturity Group Selection

Varieties of soybeans differ in maturity as much as corn varieties do, but they are classified by a different maturity system. The varieties that require the least heat to mature are placed in Group 00. In progressively warmer parts of our state, the appropriate maturities are Group 0, Group I, and Group II and early Group III. Early Group III and late mid to late Group II varieties mature dependably only in the warmer parts of central and western New York.

In the warmer regions of central and western New York, we recommend planting most of the crop with mid-Group II to early Group III varieties with a May planting date. If fields are to be planted to wheat after soybean harvest, growers should consider early Group II or late Group I varieties to insure planting wheat before mid-October. If planting occurs during the first 10 days of June, we recommend planting about 50% to early Group II and about 50% of the acreage to late Group I varieties with mid or early Group I varieties in fields that will have a subsequent wheat crop. If planting is delayed beyond June 10, we recommend planting early Group I or Group 0 varieties in central and western New York. We do not recommend planting soybeans after June 20 in these regions.

In areas of northern New York next to the lakes, we recommend planting about 50 percent of the acreage to early Group II and about 50 percent to late Group I varieties with a May planting date. Away from the lakes in northern New York, we recommend planting 70 percent of the acreage to Group I and 30 percent to Group 0 varieties. If planting is delayed until June in northern NY, we recommend planting mostly Group I varieties next to the lakes and Group 0 varieties away from the lakes. We do not recommend planting soybeans in northern NY after June 15.

6.2 Soybean Planting

The best soybean yields occur on well-drained, but not sandy, soils having a pH of 6.5 or above. The critical stage for soybean yield is in August and droughty soils that typically dry out in August will have disappointing yields. Soybeans have a very broad optimal planting date with optimum dates from about May 5-25 in the warmer regions in central and western New York. Soybeans can be successfully planted in late April or early May in these regions but final stands may be more erratic so an insecticide/fungicide seed treatment is recommended for late April and early May plantings. Mid to late Group II and early Group III varieties can be planted in these regions up until about May 20 and then just Group II varieties until June 1. If a wheat crop is to be planted after soybean harvest, then a late Group I vs. a Group II variety planted in late May will mature earlier and allow for a more timely wheat planting date. In the cooler regions in central and western New York and in Northern New York, optimum planting time is during the middle two weeks of May. Early

Group II and Group I soybean varieties should be planted at this time in these regions.

Although soybean yields decline with June plantings, high yields can still be achieved by planting early Group II or Group I varieties in central and western New York and early Group I and Group 0 varieties in Northern New York until about June 15. The earlier-maturing varieties, which tend to be short in stature, yield better at a row spacing of 15 inches or less. Soybean plantings after June 20 in central/western NY and after June 10 in NNY can be risky, even with Group 0 varieties, especially if the remaining part of the growing season is cool or if frost occurs before October 1.

It is important to place the soybean seed into the ground at a precise depth and in firm contact with the soil so choice of planting equipment is especially critical. A corn planter usually does a better job of planting than a grain drill, but soybeans typically yield about 5% less in 30-inch vs. 7.5 inch row spacing in New York even with lower final stands. In addition, modern drills have much better depth control than older grain drills.

Seeding rate depends on both row spacing and seed size. We recommend seeding rates, for seed not treated with insecticide or fungicide, of about 170,000 seeds per acre for 7.5 inch row spacing (~7.5 seeds per 3 ft.), 160,000 seeds/acre for 15-inch row spacing (about 14 seeds per 3 feet), and 150,000 plants per acre for 30-inch row spacing (~26 seeds per 3 ft.). If an insecticide/fungicide seed treatment is used, seeding rates can be reduced by 10,000 to 20,000 seeds per acre. Planting depth should be about 1.25 to 1.5 inches, depending on soil moisture conditions, and should not exceed 2 inches. Soybeans, however, can emerge reasonably well from a 2.5 inch depth, if soil crusting is not prevalent during actual emergence from the soil. Likewise, soybeans can be planted at the 1.0 inch depth, but the seed is susceptible to drying out, if conditions are dry after planting. We recommend the use of inoculum for soybean plantings in New York, especially on fields with a limited soybean history. On fields where soybeans have been grown for more than 20 years, however, inoculum may not be necessary. Likewise, the use of an insecticide/fungicide seed treatment is not necessary but can help stand establishment, especially on early-planted soybeans. Soybeans, however, can fill in the gaps very well and perfect stands are not required for maximum soybean yields.

6.3 Managing the Crop

Use soil test results to determine both lime and fertilizer requirements (see Table 6.3.1). Soybeans do not require supplemental nitrogen fertilizer if optimally fertilized for phosphorus and sulfur and at optimal pH because soybeans can fix nitrogen through a symbiotic relationship with *Bradyrhizobium* bacteria. If used, band-placed fertilizer should be at least 2 inches to the side and 2 inches below the seed. Do not place any fertilizer in contact with the

8 Appendix

8.1 Trade and Common Names of Field Crop Pesticides

Table 8.1.1 Fungicides.

Trade Name ¹	EPA Registration Number	Common Name
Absolute 500 SC	264-849	tebuconazole + trifloxystrobin
Affiance SC	10163-332	tetraconazole + azoxystrobin
Aframe Plus 2.2 SE	100-1324	propiconazole + azoxystrobin
Allegiance FL	264-935	metalaxyl
Alto 100 SL	100-1226	cyproconazole
*†Approach SC	352-840	picoxystrobin
*†Approach Prima 2.34 SC	352-883	cyproconazole + picoxystrobin
Apron XL	100-799	mefenoxam
Avaris 200 SC	100-1178-5905	azoxystrobin + propiconazole
Caramba 0.75 SL	7969-246	metconazole
Domark 230 ME	80289-7	tetraconazole
Endura 0.7 DF	7969-197	boscalid
*†Evito 480 SC	66330-64	fuoxystrobin
Fitness	34704-1031	propiconazole
Headline AMP	7969-291	pyraclostrobin + metconazole
Headline EC	7969-186	pyraclostrobin
Headline SC	7969-289	pyraclostrobin
*†Priaxor 4.17 SC	7969-311	pyraclostrobin + fluxapyroxad
Proline 480 SC	264-825	prothioconazole
PropiMax 3.6 EC	62719-346	propiconazole
Prosaro 421 SC	264-862	prothioconazole + tebuconazole
Quadris Flowable	100-1098	azoxystrobin
Quadris Top	100-1313	azoxystrobin + difenconazole
Quilt	100-1178	azoxystrobin + propiconazole
Quilt Xcel	100-1324	azoxystrobin + propiconazole
Stratego YLD	264-1093	trifloxystrobin + prothioconazole
Tilt 3.6E	100-617	propiconazole
†Topguard	67760-75	flutriafol
Topsin M 70 WP	73545-11-70506	thiophanate-methyl
TwinLine 1.75 EC	7969-247	pyraclostrobin + metconazole

*Restricted-use pesticide

†Not for use on Long Island, NY

¹Trade names are given for convenience only. No endorsement of products is intended nor is criticism of unnamed products implied.

Table 8.1.2. Insecticides.

Trade Name ¹	EPA Registration Number	Common Name
*Asana XL	352-515	esfenvalerate
*Baythroid XL	264-840	beta-cyfluthrin
*Counter	5481-545	terbufos
*Diazinon	Multiple	diazinon
Dimethoate	Multiple	dimethoate
*Force 3G	100-1075	tefluthrin

Table continues on next page.