



2025 New York and Pennsylvania Pest Management Guidelines for Grapes

Cornell Cooperative Extension



PennState Extension

These guidelines are not a substitute for pesticide labeling. Always read and understand the product label before using any pesticide.

2025 New York and Pennsylvania Pest Management Guidelines for Grapes

Authors

Bryan Brown (Cornell Integrated Pest Management Program, Cornell AgriTech, Geneva, NY; *weed management*)
Jessica Clippinger (Penn State University, North East, PA; *disease management*)
Katie Gold (School of Integrative Plant Sciences, Plant Pathology and Plant-Microbe Biology Section, Geneva, NY; *disease management*)
Bryan E. Hed (Penn State University, North East, PA; *disease management*)
Michael Helms (Cornell IPM Pesticide Safety Education, Ithaca, NY; *pesticide information*)
Greg Loeb (Department of Entomology, Geneva, NY; *entomology*)
Megan Luke (Penn State Cooperative Extension Erie County, Erie, PA; *pesticide information, pest management*)
Flor E. Acevedo (Department of Entomology, Penn State University, North East, PA; *entomology*)
Lynn Sosnoskie (School of Integrative Plant Science, Horticulture Section, Geneva, NY; *weed management*)

Special Appreciation

Special appreciation is extended to the following for their contributions to this publication:

Terence R. Bates (Lake Erie Regional Grape Program, Portland, NY); David Combs (School of Integrative Plant Sciences, Plant Pathology and Plant-Microbe Biology Section, Geneva, NY); Amara Dunn (Biocontrol Specialist, Cornell IPM, Geneva, NY); Rick Dunst (retired - Lake Erie Regional Grape Program, Portland, NY); David M. Gadoury (School of Integrative Plant Sciences, Plant Pathology and Plant-Microbe Biology Section, Geneva, NY); Heather Leach (Department of Entomology, University Park, PA); Andrew J. Landers (retired – Department of Entomology, Geneva, NY); Timothy E. Martinson (retired – Cornell Cooperative Extension, Geneva NY); Andrew J. Muza (retired – Penn State Cooperative Extension – Erie County, Erie, PA); Bruce I. Reisch, (retired – School of Integrative Plant Sciences, Horticulture Section, Geneva, NY); Michael C. Saunders (retired – Department of Entomology, University Park, PA); Andrew F. Senesac (retired – Cornell Cooperative Extension – Suffolk County, Riverhead, NY); Jody Timer (retired – Penn State University, North East, PA); Hans C. Walter-Peterson (Finger Lakes Regional Grape Program, Penn Yan, NY); Timothy H. Weigle (retired – Cornell IPM, Portland, NY); Alice V. Wise (Cornell Cooperative Extension – Suffolk County, Riverhead, NY); Wayne F. Wilcox (retired – School of Integrative Plant Sciences, Plant Pathology and Plant-Microbe Biology Section, Geneva, NY).

Abbreviations and Symbols Used in This Publication

Aacre	EC.....emulsifiable concentrate	SP.....soluble powder
AI.....active ingredient	F flowable	UDH up to day of harvest
AS.....aqueous suspension	G granular	ULV ultra-low volume
Ddust	L..... liquid	W wettable
DFdry flowable	P pellets	WDG water-dispersible granule
DGdispersible granule	PHI.....pre-harvest interval	WP wettable powder
DTH.....days to harvest	SC suspension concentrate	WSP..... water soluble packet
E.....emulsion, emulsifiable		

* Federal restricted use pesticide; may be purchased and used only by certified applicators

*^{NY} Restricted use pesticide in New York State

† Not for use in Nassau and Suffolk Counties

^ Not registered for use in New York State at press time.

Every effort has been made to provide correct, complete, and up-to-date pest management information for New York State and Pennsylvania at the time this publication was released for printing (April 2025). Changes in pesticide registrations, regulations, and guidelines occurring after publication are available in county cooperative extension offices or from Cornell IPM Pesticide Safety Education (psep.cce.cornell.edu) or the Pennsylvania Department of Agriculture's Bureau of Plant Industry (www.agriculture.state.pa.us).

Where trade names appear, no discrimination is intended, and no endorsement by Cornell Cooperative Extension or Penn State Cooperative Extension is implied.

These guidelines are not a substitute for pesticide labeling. Always read and understand the product label before using any pesticide.

The guidelines in this bulletin reflect the current (and past) authors' best efforts to interpret a complex body of scientific research, and to translate this into practical management options. Following the guidance provided in this bulletin does not assure compliance with any applicable law, rule, regulation or standard, or the achievement of particular discharge levels from agricultural land.

Cover photo by: Bryan Hed, Penn State University.

1 PESTICIDE INFORMATION.....	1
1.1 Pesticide Classification and Certification.....	1
1.2 Use Pesticides Properly	2
1.3 Pollinator Protection.....	3
1.4 State Pesticide Use Restrictions	3
1.5 Verifying Pesticide Registration	4
1.6 Check Label for Site and Pest	4
1.7 Pesticide Recordkeeping/Reporting	5
1.8 EPA Worker Protection Standard (WPS) for Agricultural Pesticides.....	6
1.9 Reduced-risk Pesticides, Minimum-risk Pesticides, and Biopesticides	6
1.10 FIFRA 2(ee) Recommendations.....	6
1.11 Endangered Species Act and Pesticides	7
2 INTRODUCTION	8
2.1 Overview	8
2.2 Specific Fungicide, Insecticide, and Herbicide Information	8
2.3 How to Develop a Vineyard Pest Management Strategy	9
2.4 Using NEWA Resources in a Vineyard Integrated Pest Management (IPM) Strategy	10
2.5 Growth Stages Critical to Grape Pest Management	11
2.6 Cornell Publications on Weeds, Insects and Diseases of Grapevines	12
2.7 Penn State Publications on Weeds, Insects, and Diseases of Grapevines	13
3 VINEYARD DISEASE MANAGEMENT	14
3.1 Pest Information – Diseases	14
Table 3.1.1 Relative disease susceptibility and sensitivity to sulfur and copper among grape varieties.....	15
Table 3.1.2 Hours of leaf wetness required for black rot infection to occur, at various temperatures following a rain.	18
Table 3.1.3 Approximate generation period for powdery mildew (time from infection by a spore until production of a colony with new spores) at different constant temperatures ^a	25
3.2 Fungicide Information.....	26
Table 3.2.1 Physical modes of action of and resistance risk of fungicides used in management of grape diseases ¹	49
Table 3.2.2 Effectiveness of fungicides for management of grape diseases ¹	51
Table 3.2.3 Relative doses of individual active ingredients provided by “combination products” when applied at label rates.....	53
4 VINEYARD INSECT & MITE MANAGEMENT.....	55
4.1 Pest Information – Insects.....	55
4.2 Insecticide Information	59
Table 4.2.1 Effectiveness of insecticides for management of grape insects and mites.	65
Table 4.2.2 Insecticides for use in New York and Pennsylvania vineyards.....	66
5 PEST MANAGEMENT SCHEDULES FOR DISEASES AND MAJOR AND MINOR INSECTS	68
5.1 Introduction.....	68
5.2 Pest Management Schedules for Diseases and Major Insects	68
5.2.1 DORMANT	68
5.2.2 DELAYED DORMANT.....	68
5.2.3 BUD SWELL.....	69
5.2.4 1-INCH SHOOT GROWTH (when the average shoot length is 1 inch)	69
5.2.5 3- to 5-INCH SHOOT GROWTH (new shoots are 3-5 inches long).....	71
5.2.6 10- to 12-INCH SHOOT GROWTH (new shoots are 10-12 inches long).....	75
5.2.7 IMMEDIATE PREBLOOM (just before blossoms open)	83
5.2.8 BLOOM (when 80-90% of the caps have fallen).....	93
5.2.9 FIRST POSTBLOOM SPRAY (10-14 days after immediate prebloom)	95
5.2.10 SECOND POSTBLOOM SPRAY (7-14 days after first postbloom spray).....	107
5.2.11 MIDSUMMER SPRAYS (July and August, as necessary)	113
5.2.12 PRE-HARVEST (Veraison to harvest period).....	124
5.2.13 POSTHARVEST	124
5.3 Pest Management Schedules for Minor and Special Insects	126
5.3.1 DORMANT TO MID-SUMMER.....	126
5.3.2 BUD SWELL TO 5-INCH SHOOT GROWTH (new shoots are 0-5 inches long)	127
5.3.3 10- TO 12-INCH SHOOT GROWTH (new shoots are 10-12 inches long).....	127

5 PEST MANAGEMENT SCHEDULES FOR DISEASES AND MAJOR AND MINOR INSECTS *(continued)*

5.3 Pest Management Schedules for Minor and Special Insects <i>(continued)</i>	
5.3.4 SUMMER TO PREHARVEST	128
5.3.5 PREHARVEST.....	129
5.3.6 PREHARVEST to POSTHARVEST.....	130

6 VINEYARD WEED MANAGEMENT132

6.1 Introduction.....	132
6.2 Resources.....	132
6.3 Effective Use of Herbicides	132
6.4 Preemergence Herbicides.....	133
6.5 Postemergence Herbicides.....	136
6.6 Herbicides for Nonbearing Vineyards	138
6.7 Specialty Use Herbicides	140

7 SPRAYER TECHNOLOGY143

7.1 Solutions for Safer Spraying – Engineering Controls	143
7.2 Preparing the Airblast Sprayer for Work	144
7.3 Sprayer Calibration	145
7.4 Spray Nozzle Types	148
7.5 Selecting Nozzles from the Nozzle Catalogue – Airblast Sprayers	148
7.6 Selecting Nozzles from the Nozzle Catalogue –Boom Sprayers	150
7.7 Equipment for Weed Control in Vineyards	151
7.8 Drift Management Strategies	152
7.9 Reducing Drift from Airblast Sprayers.....	153
7.10 Going Spraying!.....	154
7.11. Cleaning Sprayers	157
7.12 Spraying Small Vineyards	160
7.13 Application Equipment Distance Learning.....	163

8 PESTICIDES FOR NEW YORK AND PENNSYLVANIA VINEYARDS164

8.1 Herbicides.....	164
8.2 Insecticides and acaricides.....	165
8.3 Fungicides.....	166

1 Pesticide Information

1.1 Pesticide Classification and Certification

Pesticides can be classified as general use or restricted use. **General use pesticides** may be purchased and used by anyone. **Restricted use pesticides** can only be purchased and used by a certified applicator or used by someone under a certified applicator's supervision. In some cases, the pesticide label may limit use of a restricted use pesticide to only a certified applicator.

1.1.1 Certification in New York State

Private applicators use or supervise the use of pesticides to produce agricultural commodities or forest crops on land owned or rented by the private applicator or their employer. If a private applicator wants to use or supervise the use of restricted use pesticides, they need to be a **certified private applicator**. Certified private applicators are also allowed to purchase restricted use pesticides. Certification is not needed if a private applicator uses general use pesticides.

In New York State, a **certified commercial applicator**, **certified commercial technician**, or **commercial apprentice** working under the supervision of a certified commercial applicator is allowed to apply any type of pesticide on property that is not a private application (described above) or is a residential application. (A residential application is the use of general use pesticides on property owned or rented by the applicator, excluding establishments selling or processing food and residential structures other than where the applicator lives.) Certified commercial applicators are allowed to purchase restricted use pesticides.

Information on pesticide certification and classification is available from your Cornell Cooperative Extension office (cce.cornell.edu/localoffices), regional NYSDEC pesticide specialist (dec.ny.gov/about/contact-us/statewide-office-information), the Pesticide Applicator Training Manuals (www.cornellstore.com/books/cornell-cooperative-ext-pmep-manuals), or the Cornell Pesticide Safety Education Program (psep.cce.cornell.edu).

1.1.2 Certification and Business Requirements in Pennsylvania

In Pennsylvania, definitions of private applicator and commercial or public applicator are similar as noted above. Additionally, pesticides may be applied by registered technicians and/or noncertified applicators under the direction of a licensed applicator.

Private Applicators – Private applicators are an applicator who intends to purchase and/or apply restricted-use pesticides for the purpose of producing an agricultural

commodity on land that is owned or rented by that person or their employer.

Commercial and Public Applicator - There are 25 categories of commercial and public applicators. To be eligible for certification, a person must pass two written examinations (core and category) and be employed by a pesticide application business. Once exams have been successfully passed, that person has 12 months to apply for certification with the Pennsylvania Department of Agriculture.

Application Business – The Pennsylvania Department of Agriculture (PDA) defines a pesticide application business as a government entity or commercial establishment engaged in the application of pesticides and meeting the definition of a commercial or public applicator. This includes individual operators that meet the definition of a commercial or public applicator.

Businesses must be licensed to apply general or restricted-use pesticides. A license must be obtained indicating the certification category in which the business makes applications. The business must, at all times, employ an applicator certified for each category in which it intends to make a pesticide application.

Businesses must meet the requirements for comprehensive general liability insurance coverage for pesticide applications. A minimum of \$200,000 coverage is required. This must include \$100,000 coverage for bodily injury and \$100,000 coverage for property damage per occurrence. Coverage must include a statement that pesticide applications are included, or endorsement #CG26160194 or #CG26160798.

Registered technicians – Registered technicians are those applicators who have received training as required under subsection 128.51 of the Pennsylvania Pesticide Rules and Regulations. These applicators have completed minimum training requirements under the direction of an applicator with at least one year's certification in the category for which the technician is being trained. They must also be under the supervision of a certified applicator who can be on site within five hours if necessary. Registered technicians must be re-registered and receive updated training on an annual basis.

Noncertified Applicators – Applications may be made by an individual who is neither certified nor a registered technician if and only if that individual is under the direct supervision of a certified applicator who is physically present and within sight of the application being made.

2 Introduction

2.1 Overview

How do you decide which pest management strategy is right for you? The variety of choices can be daunting: integrated pest management, sustainable, integrated crop management, organic, best management practices, biodynamics, or traditional, just to name a few. However, what it boils down to is that it doesn't really matter what you call your pest management strategy. Consider which practices work best for you in a particular vineyard. Take the best tools from each of the different programs to provide you with the management strategy that fits your needs and pocketbook.

The one component that every successful pest management strategy shares is information. By gathering and compiling the right information, you will be able to make the best choices in putting together your vineyard pest management strategy.

2.2 Specific Fungicide, Insecticide, and Herbicide Information

The following materials have been registered for the control of certain insects, diseases, and weeds affecting grapes. The trade name and (common) name are given, as well as selected uses for each material. The signal word associated with each pesticide indicates its relative level of toxicity. High toxicity indicates that small quantities of the chemical may cause serious illness or death.

2.2.1 Pesticide Toxicity

The toxicity of a chemical typically is measured with a Lethal Dose 50 (LD50) value. This value is the dosage necessary to kill 50 percent of a laboratory population of test animals (rats, mice, or rabbits). These toxicity values may be expressed in terms of a single dosage in milligrams per kilogram of body weight. LD50 values are useful in comparing different pesticides, as the degree of hazard to a person handling a pesticide is directly related to toxicity.

The following classification was established to aid users of pesticide chemicals:

High toxicity: Acute oral LD50 from a trace to 50 mg/kg. From 4 to 100 drops (1 teaspoon) of technical pesticide may be lethal to a 150-pound person. Label must carry signal word DANGER. A skull and crossbones on the label indicates poison.

Moderate toxicity: Acute oral LD50 from 50 to 500 mg/kg. From 1 teaspoon to 1 ounce of technical pesticide may be lethal to a 150-pound person. Label must carry signal word WARNING.

Low toxicity: Acute oral LD50 greater than 500 mg/kg. From 1 ounce to 1 pint of technical pesticide may be lethal to a 150-pound person. Label must carry signal word CAUTION.

2.2.2 Pesticide Compatibility

Some pesticides are unstable or incompatible under alkaline conditions. Well or pond water used to fill sprayers can be alkaline, and pH should be tested and adjusted if over 7.0. Do not apply tank-mix combinations unless your previous experience indicates the mixture is effective and will not result in application problems or plant injury. If tank-mix compatibilities are unknown, the mixture should be combined in the proper proportions using a jar test. CAUTION: Fixed copper formulations and lime should not be used with Captan, Imidan, or Sevin.

2.2.3 Adjuvants with Fungicides and Insecticides

The addition of adjuvants (spreader-stickers, penetrants, etc.) to spray mixtures of fungicides and insecticides is not recommended unless suggested on the pesticide label or supported by reliable data. Many fungicide and insecticide formulations already include an adjuvant; thus, addition of another adjuvant is often counterproductive or a waste of money.

2.2.4 Pesticide Resistance Management

The ability of various fungicides, insecticides, and herbicides to control specific grape pests has been reduced (and, in some cases, eliminated) because pests developed resistance to these materials. Resistance occurs when some individual pests survive treatment with the pesticide. They multiply and pass this resistance on to their young. Because individuals that are controlled by the chemical die, or reproduce poorly, over time the population becomes dominated by individuals that are resistant to it. This process is favored by a number of factors, the most common of which are:

1. Multiple applications of a single product or class of compounds (strobilurin fungicides, organophosphate insecticides, etc.);
2. An exclusive reliance on a single product (or class) to manage the pest;
3. Repeated treatments of large pest populations with the products (e.g., "rescue" treatments); and, in some cases
4. Attempts to manage the pests with very low rates of the products (either intentionally or due to poor spray coverage).

Although it's not possible to guarantee that resistance will never develop to any specific pesticide, ALL growers should follow a few simple rules to minimize the probability of this occurring. These rules are particularly

2.5 Growth Stages Critical to Grape Pest Management (*continued*)



new shoots, 1/2 – to 1-inch stage



new shoots, 3- to 5-inch stage



new shoots, 10- to 12-inch stage



trace bloom



postbloom

2.6 Cornell Publications on Weeds, Insects and Diseases of Grapevines

2.6.1 Fact Sheets

Cornell vineyard weed management fact sheets are available free upon request from the Bulletin Room, New York State Agricultural Experiment Station, Geneva, NY 14456.

Pool, R.M., R.M. Dunst, and A.F. Senesac. 1990. Managing weeds in New York vineyards. I. Choosing a weed management program. Cornell Grape Facts I-1.

Senesac, A.F., R.M. Dunst, and R.M. Pool. 1990. Managing weeds in New York vineyards. II. Chemical control of vineyard weeds. Cornell GrapeFacts I-2.

Dunst, R.M., A.F. Senesac, and R.M. Pool. 1990. Managing weeds in New York vineyards. III. Pre-emergence herbicides. Cornell GrapeFacts I-3.

Dunst, R.M., A.F. Senesac, and R.M. Pool. 1990. Managing weeds in New York vineyards. IV. Post-emergence herbicides. Cornell GrapeFacts I-4.

3 Vineyard Disease Management

3.1 Pest Information – Diseases

Several important insects and diseases that occur in New York and Pennsylvania vineyards are described below to help growers manage these pests with practices and pesticides appropriate for their production systems.

ANGULAR LEAF SCORCH was first described in 1985.

Symptoms of this fungal disease are similar to those of rotbrenner, a disease of grapevines found in the cool grape-growing regions of Europe, which is caused by a very closely related fungus. Angular leaf scorch occurs sporadically and is most likely to become a problem in years when high rainfall occurs between bud break and early summer, especially if this should happen in consecutive years. Riesling appears to be a particularly susceptible cultivar.

Disease symptoms occur mainly on the leaves and first appear as faint chlorotic spots. As these lesions grow larger, they change from yellow to reddish-brown and the margin often becomes sharply defined (depending on the cultivar, the margin may be yellow, red, or absent). Lesions are confined by major veins, becoming "angular" or wedge-shaped. They eventually kill the infected tissue, sometimes crossing the major veins in the process and often causing diseased leaves to fall prematurely.

The fungus survives winter in infected leaves on the vineyard floor. Mature spores are ready for discharge in spring when grape buds begin to grow. During rainfall, spores are released into the air from fruiting structures, and susceptible tissue is infected.

Cultural practices that increase air circulation through the canopy can shorten periods of leaf wetness that favor disease development. Destruction of leaf litter by cultivation, before bud break, can also reduce disease pressure. Where needed, effective fungicides applied from the 3-inch stage and continuing through fruit set will provide significant control. Although there are no specific labels for the control of this disease, mancozeb products (e.g., Dithane, Penncozeb) applied according to label directions to control *Phomopsis*, black rot, and downy mildew also have provided good control of angular leaf scorch in commercial experience. In Europe, the strobilurin fungicides (Abound, Flint, Pristine, *^{NY}Sovran) have provided good control of the closely related disease, rotbrenner. However, experience with angular leaf scorch is lacking. The DMI fungicide, difenoconazole (Revus Top, Quadris Top; CAUTION: Quadris Top causes injury to Concord and some other cultivars) is labeled for control of rotbrenner and should provide significant post-infection control of this disease in addition to moderate protective activity (excellent protective activity for Quadris Top). See Table 3.1.1 for varietal susceptibility to this disease.

ANTHRACNOSE is a disease that occurs most commonly in years that are wet during the first half of the growing season, with damage typically limited to a few highly susceptible cultivars. In NY/PA, most outbreaks historically occurred on Vidal Blanc and a few seedless table grape varieties, especially Reliance. In recent years, however, there have been regular outbreaks on some of the newer cold-hardy cultivars that are gaining in popularity and expanding the geographical range of grape production. Marquette appears to be particularly susceptible, although Frontenac and La Crescent also have been affected. Some older cold-hardy cultivars (Edelweiss, Esprit, Brianna, St. Pepin, and Swenson White) also can be problematic. In some Midwestern states, Concord, Catawba, and Leon Millot have been reported as encountering problems, although such occurrences are rare in NY and PA. Symptoms occur on leaves, green shoots, and clusters. On leaves, numerous small, circular brown spots appear which later turn gray in the center and develop dark brown to black margins. In severe attacks, lesions may coalesce and cause large dead zones, distortion of the leaf blade, and eventually death of the entire leaf. Infected shoots develop dark, noticeably sunken lesions, typically on the first several internodes near the base of the new shoot. These lesions resemble the internode lesions typical of *Phomopsis* cane and leaf spot but they usually are more aggressive, expanding farther along the shoot and deeper into its center than those caused by *Phomopsis*. On berries, spots approximately 0.25-in in diameter develop, with whitish-gray centers surrounded by reddish brown to black margins, sometimes producing an appearance that superficially resembles a bird's eye. Severely affected berries may shrivel and dry into mummies.

The fungus overwinters primarily on infected canes, although the previous year's berries can also be a source. In spring, spores are produced from the fungal structures on these sources and are dispersed by splashing raindrops to young, susceptible tissues, where they cause infection if wetness persists for a sufficient length of time. Temperatures in the mid-70s to mid-80s Fahrenheit (25-30°C) are optimal and require only 3 to 4 hours of leaf wetness for infection to occur. However, infection can also occur across a much wider range of temperatures, including those that typically prevail during the early growing season in upstate NY, if it remains wet for long enough. Additional spores, which also are splash dispersed, are produced from new infections, and these can rapidly spread the disease through multiple repeating cycles of new infection and additional spore production. Hence, outbreaks occur most frequently in years with multiple rain events early and mid-season. Young tissues are most susceptible, becoming resistant as they are mature; for example, berries become relatively resistant by about 7 weeks post-bloom.

Table 3.1.1 Relative disease susceptibility and sensitivity to sulfur and copper among grape varieties¹.

Variety	Disease susceptibility or chemical sensitivity ^a									
	BR	DM	PM	Bot	Phom	Eu	CG	ALS	S ^c	C ^d
Sauvignon blanc	+++	+++	+++	+++	?	+++	+++	?	No	+
Seyval	++	++	+++	+++	+++	+	++	++	No	+
Steuben	++	+	+	+	?	?	+	++	Yes	?
Traminette	+++	++	+	++	?	?	+	?	No	?
Valvin Muscat (NY62.0122.01)	++	+	++	+	?	?	+	?	No	?
Vanessa	+++	++	++	+	+	?	+	?	No	?
Ventura	++	++	++	+	+	?	+	+++	No	?
Vidal 256	+	++	++	+	+	+	++	+	No	+
Vignoles	+	++	+	+++	+++	++	++	++	No	?
Key:	+ Slightly susceptible or sensitive			No = Not sensitive						
	++ Moderately susceptible or sensitive			? = Relative susceptibility or sensitivity not established						
	+++ Highly susceptible or sensitive									

¹The relative ratings in this chart apply to an average growing season in NY and PA. Under conditions favorable for disease development, any given variety may be more severely affected.

a. BR=Black rot, DM=Downy mildew, PM=Powdery mildew, Bot=Botrytis, Phom=Phomopsis, Eu=Eutypa, CG=Crown gall, ALS=Angular Leaf Scorch, S=Sulfur, C=Copper

b. Berries only weakly susceptible

c. Slight to moderate sulfur injury may occur even on tolerant varieties when temperatures are 85F or higher during or immediately following the application

d. Copper is most likely to cause injury when applied under slow-drying conditions (cool or very humid).

Diseased canes should be pruned during the dormant season and removed from the vineyard or destroyed. If numerous infected berries remain on the vineyard floor, the spores originating from them can be largely neutralized by covering the berries with soil through cultivation or, if practical, covering them with mulch. Early-season sprays of mancozeb, captan, or ziram targeted against Phomopsis also provide significant control of anthracnose, although this latter disease is not listed as a target on most labels. Some DMI (Group 3) fungicides, e.g., difenoconazole (Revus Top, Quadris Top, and Inspire Super), myclobutanil (Rally), and tetraconazole (Mettle) are specifically labeled for anthracnose control. A “delayed dormant” application of lime sulfur can be useful in vineyards where the disease has become established and problematic to control and/or in “organic” vineyards where traditional fungicides are not used. This treatment limits the production of infectious spores from overwintered cankers but does not protect new growth from any spores that survive the application or are produced after its effects wear off.

BITTER ROT and RIPE ROT are two diseases that are common in states to the south of PA where temperatures are consistently warmer, although they also occur sporadically in NY and PA, especially in the southern regions of our two states. The two diseases are caused by different fungi, but both have similar biologies and respond to the same management practices. Bitter rot is active at somewhat lower temperatures than ripe rot and is the more regular threat in the mid-Atlantic region, although their geographical ranges overlap. In NY, bitter rot has been most problematic on Long Island, primarily

in years with wet summer and pre-harvest periods (particularly on Chardonnay). It has also been problematic in southeast PA in some years, and both diseases occur further north on occasion. Usually, bitter rot symptoms first develop after veraison, when the causal fungus moves into the berry from the infected berry stem and turns the diseased portion brown (on white varieties) or a dull purple. Once the berry is completely rotted, it becomes covered with numerous prominently raised black pustules (the fungal fruiting bodies, called acervuli). Within a few days, diseased berries soften and may drop; others shrivel into mummies that resemble those caused by black rot and Phomopsis. This late onset of symptom appearance is one feature that distinguishes bitter rot from black rot (virtually all black rot infections should be apparent by veraison); others include (i) the appearance of the fungal fruiting bodies on infected fruit (those of black rot are relatively small, round and uniform in size, whereas those of bitter rot are larger and much more pronounced, providing a bumpy texture to the berry); (ii) the tendency of black-rot infected fruit to remain attached to the berry stem when pulled gently, whereas those with bitter rot are more easily detached; and (iii) the tendency of fruit infected with bitter rot to leave hands sooty black if handled when wet, whereas those infected with black rot will leave hands clean. Berries infected with Phomopsis also tend to first appear during the preharvest period but are detached very easily when pulled, and the black fruiting bodies produced on rotten berries are relatively few and inconspicuous. Also, significant Phomopsis fruit rot typically occurs in association with significant visible infections of the cluster stem and of the shoots and

particularly positive impact on the control of cluster infections the following season.

For effective management of powdery mildew, sprays may be required as early as 3- to 5-inch shoot growth on *V. vinifera* and highly susceptible hybrid cultivars, depending on rainfall and temperature in the early season. Note that cold nights near 40°F and below can significantly retard disease development, so spring seasons with overcast skies that minimize such episodes are at greater risk than those with a number of clear, cool nights.

Sprays may need to begin by the 10-inch shoot stage on Concord or other moderately susceptible cultivars, and should not be delayed beyond the immediate prebloom stage on any but the most highly resistant cultivars; sprays should be initiated immediately if the disease is observed before then. To protect against powdery mildew developing on the fruit, management programs should be at their peak intensity from just before bloom through fruit set or shortly thereafter, emphasizing the use of highly effective fungicides, full rates, appropriate spray intervals, and superior spray coverage (every row, proper speed, sufficient water volume). Spray programs may need to be especially “tight” when days and nights remain warm (see Table 3.1.3) and/or weather is cloudy and damp during this period. Protection of Concord berries is not required after fruit reach 0.25-inch in diameter, although continued foliar protection is likely to be beneficial under high-crop or poor-ripening conditions. For *V. vinifera* and susceptible hybrids, it is important to maintain excellent protection of the clusters through the bunch closure period, since powdery mildew infections at this time can promote the later development of bunch rots and/or wine spoilage microorganisms. Maintenance programs to protect foliage throughout the summer are necessary for attaining maximum fruit and vine quality on *V. vinifera* and susceptible hybrid cultivars. Good control of foliar infections through Labor Day will contribute significantly to the control of fruit infections the following year, since it will limit the overwintering inoculum needed to produce them.

RIPE ROT (see “Bitter rot”)

SOUR ROT develops on injured berries when the weather is wet during the pre-harvest period, particularly if it is relatively warm (above 60°F, with 70s being optimum for disease development) for a significant length of time then. The characteristic vinegar smell from which the disease gets its name results from the activity of a specific group of bacteria, which oxidize the ethanol that wild yeasts produce from the juice of the injured berries. Such berries may also show signs of various “secondary” fungi that colonize decaying fruit, and may further be colonized by certain other yeasts that produce a compound (ethyl acetate) that smells like nail polish remover, which detracts from the quality of finished

wines. Numerous fruit flies (sometimes called vinegar flies) – i.e., *Drosophila* spp. – also swarm around decaying berries and multiply by laying their eggs within them. These insects play a central role in the development of the disease and furthermore spread the decay organisms throughout the vineyard once some clusters begin to break down. The sour rot organisms also are spread as diseased berries drip contaminated juice onto nearby healthy berries, which in turn become infected through any wounds that might be available. Rain cracks, bird and insect damage, Botrytis or powdery mildew infections, and mechanical injuries caused as swelling berries pull away from their stems in tightly compacted bunches are among the many injury sites in which sour rot can become established.

Sour rot can increase very rapidly during the pre-harvest period, and it long has been thought to be almost impossible to stop the disease once it has become established if warm and wet conditions persist. However, recent research suggests that disease development can be reduced through a combination of insecticide sprays to control *Drosophila* fly populations and registered broad-spectrum sterilants (e.g., Oxidate) to limit microbial populations on the berries. This program is most effective if started around 15°Brix, as berries become susceptible and before symptoms develop, but it also can reduce further spread if initiated immediately after the disease is noticed through scouting activities. Limiting various causes of injury will also greatly reduce sour rot development. Open canopies and leaf removal around clusters, which allow them to dry once rains have stopped, can lower disease severity significantly. As with Botrytis, practices that loosen cluster compaction can have a major effect on limiting sour rot development; however, most of these practices remain experimental or difficult to employ.

3.2 Fungicide Information

ABOUND (azoxystrobin) – read the label

Signal word: CAUTION

Medical emergency: (800) 888-8372

Chemical/fungicide family: strobilurin [QoI]

Resistance Group Number: 11

Resistance risk: high

Physical mode of action: protectant, variable but limited post-infection, antisporeulant

Selected uses: powdery mildew, downy mildew, black rot; modest suppression of Phomopsis and Botrytis

Comments: Azoxystrobin is the active ingredient in Abound. It also is formulated with DMI fungicides, to make two different combination products, Quadris Top (in combination with difenoconazole) and †Topguard EQ (in combination with flutriafol), and with mancozeb to make *^{NY}Dexter Max. Abound was the first registered compound in the strobilurin (or “QoI”) group of

***NY†ZAMPRO 4.4SC** (ametotradin + dimethomorph) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (800) 832-4357

Chemical/fungicide family: triazolo-pyrimidylamine (QxI) + carboxylic acid amide (CAA)

Resistance risk: moderate

Resistance group: 45, 40

Physical mode of action: protectant, some post-infection, anti-sporulant

Selected use: downy mildew (only)

Comments: *NY†Zampro is a new downy mildew-specific fungicide. It contains two active ingredients, (i) ametototradin, which represents a new class of chemistry, with no other similar compounds yet on the market; and (ii) dimethomorph, an older Group 40 product in the same class as Revus (the downy mildew component of Revus Top). It has provided excellent control of downy mildew in several NY trials and elsewhere. Its physical mode of action is not well characterized, but it appears to have good protective activity; significant but limited post-infection activity; and anti-sporulant activity. *NY†Zampro has a 12 hr REI and a 14-day PHI.

RESISTANCE WARNING: Downy mildew resistance to Group 40 fungicides has been found in three vineyards in Virginia and one in North Carolina (Feng and Baudoin, 2018). **Downy mildew resistance to Group 40 fungicides has been found in high amount throughout the Lake Erie grape growing region (PA and NY) in 2023 and 2024. FRAC 40 fungicides should not be solely relied upon for DM control in this region.** No

more than two applications of *NY†Zampro and Revus (or Revus Top) **combined**, and no sequential applications of a Group 40 products are recommended. Zampro should still control downy mildew if FRAC 40 resistance is present due to the FRAC 45 component, but also prone to resistance development and should be limited in use to twice a season, applied in rotation with unrelated FRAC group products, and avoid back-to-back applications.

ZIRAM (ziram) – read the label

Signal word: DANGER

Medical emergency: (866) 673-6671

Chemical/fungicide family: dimethyldithiocarbamate

Resistance Group Number: N/A

Resistance risk: low

Physical mode of action: protectant

Selected uses: black rot, Phomopsis; some downy mildew

Comments: Do not enter the treated area within 48 hours after application unless proper protective clothing is worn. Ziram is similar to mancozeb products in its activity against black rot and Phomopsis, but is less effective than mancozeb against downy mildew. Ziram can be used from the 6-inch shoot growth stage until 21 days before harvest (maximum 24 lb/A/year). It is especially useful as a postbloom substitute for mancozeb where processor restrictions prohibit use of the latter. **Note: Ziram has been canceled in grapes by the EPA, however, the 12 month grace period has not started as of printing. Therefore, Ziram can continue to be used through the 2025 season.**

Table 3.2.1 Physical modes of action of and resistance risk of fungicides used in management of grape diseases¹.

Fungicide	Protectant ^a	Post-infection ^b	Anti-sporulant ^c	Eradicant ^d	Resistance risk	Resistance group
ametotradin + dimethomorph (*NY†Zampro)	+	+	+	-	M	45, 40
azoxystrobin (Abound, Quadris Top, †Topguard EQ, *NYDexter Max) ^e	+	+/- ^f	+	-	H	11
<i>Bacillus amyloliquefaciens</i> F727 (Stargus)	+	-	-	-	L	BM02
<i>Bacillus mycoides</i> isolate J (LifegardWG)	+	-	-	-	L	N/A ^j
<i>Bacillus subtilis</i> strain AFS032321 (Theia)	+	-	-	-	L	BM02
benzovindiflupyr (*NYAprovia, *NYAprovia Top)	+	+	+	-	M	7
boscalid (Endura)	+	+	+	-	M/H	7
boscalid + pyraclostrobin (Pristine)	+	+/- ^g	+	-	M/H	7, 11
captan (Captan)	+	-	-	-	L	N/A ^j
Cerevisane (Romeo)	+	-	-	-	L	BM02
copper (several formulations)	+	- ⁱ	-	-	L	N/A ^j
cyazofamid (Ranman)	+	+/- [?]	+	-	H?	21
cyprodinil (Vangard, †Switch)	+	+	+	-	M/H	9
cyflufenamid (Torino)	+	+	+	-	M/H?	U6

Table continued on next page.

4 Vineyard Insect & Mite Management

4.1 Pest Information – Insects

BANDED GRAPE BUG is a sporadic pest of grapes in the Finger Lakes and Lake Erie regions and does not require treatment in most years. Nymphs of this insect emerge in the spring and feed on flowers and young berries, using their sucking and piercing mouth parts. The nymphs range in size from 1/8- to 1/2-inch in length, depending on the stage. Injury by small nymphs, occurring between 3- to 5-inch shoot growth (around May 15) and early June, results in floret drop, reduced berry set, and fewer clusters. Subsequent feeding by larger nymphs and adults does not affect cluster development. Economic injury can occur when more than 1 nymph per 10 shoots are present. This injury only occurs in the prebloom stages. Subsequent feeding by nymphs does not reduce berry set. Adults appear to be predaceous and do not cause injury to berries. Look for nymphs on grape clusters and shoot tips prior to the bloom period. They can be recognized by their long, banded antennae.

BROWN MARMORATED STINK BUG (BMSB) is a new invasive species in New York that may present problems for grapes. This stink bug, originally from Asia, was first observed in PA and has spread to many regions and has become particularly abundant in the mid-Atlantic states. It is present in NY and PA grape-growing regions, although at this time at relatively low numbers. BMSB uses its sucking mouthparts to feed on reproductive structures of many different crop plants, including grapes. At high densities, damage can be extensive. BMSB also produces strong odors that have the potential of tainting grape juice. Recent research indicates the offending compounds are not very stable and break down during fermentation. Even without fermentation, odors are relatively unstable and may not be of significant concern except when consumed close to harvest.

CLIMBING CUTWORMS are known to feed on grapes. Larvae hide in the soil litter below the grape trellis and climb onto vines on warm nights to feed on developing primary buds. Only during bud swell are cutworms able to inflict serious damage to a vineyard. To examine vines for cutworms, search under the bark and in the soil litter beneath a vine with damaged buds, or search the vine with a flashlight after dark.

EUROPEAN CORN BORER is an important lepidopteran pest of corn, but it is also known to feed on over 200 other plant species, including grapes. Corn borer problems are rare, but under some circumstances, may require management. They are usually found in *Vinifera* varieties, especially vines with excessive foliage or where vineyards are weedy or surrounded by corn, sorghum, Sudan grass, or related crops. Young vineyards or nursery stock may be more seriously affected by borer

injury than mature vines. The larvae vary in color, ranging from creamy to light gray to faint pink, with very small, round, dark brown spots on each segment and a dark-colored head capsule. After initially feeding on young leaves, larvae bore into canes. This weakens or kills shoots, especially when the larvae enter the middle or lower sections. Adult moths are a creamy yellowish-brown and approximately one inch long. Eggs are white and laid in masses resembling overlapping fish scales on the underside of leaves. Egg laying can occur in late May, late June to early July, or early August, depending on the genetic race of corn borer present. See the section on pest management schedules for minor insects (4.3) for pesticide recommendations and other comments.

EUROPEAN RED MITES are spider mites. Adult mites are small, dark red, and have eight legs. When viewed with a hand lens, the mites appear hairy because they have white spines called “setae.” Nymphs range in color from pale to dark orange. Both adults and nymphs pierce the leaf cells and extract plant juices. This leads to the characteristic bronze coloration, which impairs the photosynthetic capacity of the leaf. Two-spotted spider mites are often found in mixed populations with European red mites. Two-spotted spider mites are light in color with two black spots on their backs. *Vinifera* and French hybrid varieties appear to be the most susceptible to infestations, although native varieties can also develop large densities under some conditions. Mites may be found on the upper or lower leaf surface. Four to nine generations occur in a season. Susceptible vineyards in production areas prone to damaging infestations should be monitored, starting at the bud break stage, for presence of this pest. Although problems can develop at any time after bud break, pay particular attention to the 1- to 4-inch growth stage and the postbloom period, especially after early July. Given a head start, the vine can tolerate a fair amount of feeding damage on lower leaves. Heavy mite infestations early in the season can cause stunted, chlorotic shoots with small leaves and pinpoint necrotic areas on leaves. Later in the season, as shoot growth rate declines and the vine allocates more resources to fruit, mites may also have an increased capacity to cause damage. Infestations can be severe on Long Island and in southeastern Pennsylvania vineyards. Serious infestations in the Finger Lakes region have occurred more frequently in recent years. Problems with spider mites in the Lake Erie region are uncommon. Predatory mites, when present in the vineyard at sufficient densities, can provide excellent biological control of spider mites. Recent research indicates that frequent use of mancozeb fungicides reduces predatory mite populations, although mancozeb use does not necessarily lead to mite problems.

GRAPE BERRY MOTH is one of the most serious insect pests affecting grapes in New York and Pennsylvania.

mouth parts to feed on leaf tissue. When populations are high, during the early part of the season, their feeding activity can result in small, deformed leaves and stunted shoots. Later in the season, the vines are much better able to tolerate thrips feeding. It is rarely necessary to treat this pest. Aurore and DeChaunac varieties appear to be most susceptible to shoot stunting. Concord and other native varieties tolerate feeding with no apparent injury.

TUMID GALLMAKER is a small (1/10-inch) brown to reddish fly with plume-like antennae. From early May to mid-September, it lays its eggs in masses between developing tissues at the bud or shoot tips. The larvae cause injury to grapes. After hatching, they bore into vine tissue and cause a round, reddish gall to form. These galls can develop on leaf tissue or petioles, where they probably do little actual damage to the vine, or in grape clusters, where there is more concern about economic injury. Hence, the greatest concern for this pest is in the early part of the season. Tumid gallmaker is generally not as prevalent in the western and central grape-growing regions as in the southeastern areas. Aurore and Rougeon appear to be particularly susceptible.

YELLOWJACKETS, OTHER SOCIAL WASPS

Yellowjackets and some other types of social wasps can be problematic at or near harvest in vineyards. Although predaceous during most of the year and therefore, beneficial, near harvest yellowjackets and other social wasps (paper wasps, hornets) are attracted into vineyards in search of berry juice (sugar). They will feed on previously damaged berries and for some species of yellowjackets, will also directly feed on and damage intact berries, opening them up to bacteria, yeast and vinegar flies, key contributors of sour rot. Yellowjackets and other social wasps can also create significant nuisance issues. Management options are limited. No insecticides are labeled to control yellowjackets in vineyards, although some of the insecticides labeled for use against spotted wing drosophila, such as *Mustang Maxx and Delegate, may provide some short-term knock down of foraging yellowjackets. Finding nests earlier in the season and treating with labeled insecticides may also help reduce overall populations closer to harvest. And there is some research showing that using traps with an attractive lure and killing agent can help reduce yellowjacket populations. Finally, during the 2023 and 2024 field seasons we tested the use of side netting with smaller mesh size (3 X 5 mm) and found a significant reduction in yellowjacket feeding damage compared to vines without netting. Overall, however, more research is needed geared toward developing management tactics for yellowjackets.

4.2 Insecticide Information

ACRAMITE (bifenazate) – read the label

Signal word: CAUTION

Medical emergency: (866) 673-6671

Selected use: spider mites

Comments: Do not apply within 14 days of harvest. REI is 5 days for tying, turning, and girdling of table grapes and 12 hrs for all other activities. Both European red mite and two-spotted mite are listed for grapes on the current label. Acramite limited to one application per season.

***NY†ACTARA** (thiamethoxam) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (800) 888-8372

Selected use: leafhoppers, mealybugs, Japanese beetle

Comments: REI = 12 hrs, DTH = 5 days. Maximum *NY†Actara allowed per growing season is 7 oz/A with a minimum interval between applications of 14 days. Not registered for use in Nassau and Suffolk Counties in New York State.

***NY ADMIRE PRO** (imidacloprid) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (800) 334-7577

Selected use: mealybugs, leafhoppers, grape phylloxera

Comments: REI = 12 hrs, DTH = 30 days (**soil**) & 0 days (**foliar**) *NY Admire Pro can now be applied both to soil, where it is systemic throughout the vine, and on foliage, where it shows translaminar activity. When applied to soil, *NY Admire Pro works best when applied through a drip system, although it can be applied as a subsurface side-dress. Maximum allowed per season is 14.0 fluid ounces per acre for **soil treatment** and 2.8 fluid ounces per acre for **foliar treatment**. [*NY Alias 4F is a generic of soil applied imidacloprid that is also labeled for foliar application.]

***AGRI-MEK** (abamectin) – RESTRICTED-USE PESTICIDE – read the label

Signal word: WARNING

Medical emergency: (800) 888-8372

Selected use: two-spotted spider mite

Comments: REI = 12 hrs, DTH = 28 days. Product must be applied in combination with nonionic surfactant. Do not make more than two applications per season. Amount of product per growing season cannot exceed 32 fl oz/A of 7.0 fl oz *Agri-Mek SC. Ground application only. Currently, European red mite is not on the label..

***NY†ALTUS** (flupyradifurone) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (800) 334-7577

Selected uses: Leafhoppers, some other sucking insects.

Comments REI = 4 hrs; DTH = 0 days. This insecticide belongs to a new class of chemicals known as

5 Pest Management Schedules for Diseases and Major and Minor Insects

5.1 Introduction

This section provides guidelines pertaining to management programs for control of diseases and major insects in vineyards of New York and Pennsylvania. Although this section is organized along a phenological schedule to reflect important events during the growing season, there is no implication that every spray listed will be necessary. Rather, this is a schedule of the various times when individual diseases and insects might require that sprays be integrated into a management program; refer to the notes to help determine which sprays are generally necessary and

which ones apply only to certain conditions. Refer to the pictures in the front of this publication for help in identifying critical growth stages during the season. Note comments in right-hand column address precautions or considerations necessary for use of particular methods or materials. Be especially alert to the notations that certain chemicals may not be approved for your state or for certain growing areas within a state. See section 5.3 for pest management schedule for minor or special insect pests such as spotted lanternfly.

5.2 Pest Management Schedules for Diseases and Major Insects

Pest(s)	Materials	Rate per Acre	Comments
5.2.1 DORMANT			
Canker diseases (Eutypa, Botryosphaeria)	Mettle 1ME	5 fl oz	Mettle is labeled in all states for spray application in 25 to 50 gpa within 24 hr after pruning, with a 12-hr REI. Consult the label for further use directions. No trials have been conducted in NY or PA to evaluate the efficacy of Mettle for this purpose.
5.2.2 DELAYED DORMANT			
Soft scale insects and mealybugs	petroleum oil	2.5%	Apply early in the spring at bud swell but before any leaf tissue is exposed. Apply in 250 to 300 gallons of water. Thorough coverage is essential for good results. Avoid use with captan or sulfur due to phytotoxicity. Also avoid use within 24 hours before or after freezing temperatures. Field data indicate only short-term benefits for reducing mealybug populations. Oil more effective against soft scale.
	OR Knack	16 fl oz	Only labeled for lecanium scale. Can use oil with Knack at this time (delayed dormant). Not to exceed 32 fl oz/A for season.
Anthracnose, black rot, Phomopsis, powdery mildew	* ^{NY} Miller Lime Sulfur	1 gal/10 gal water	This spray is most likely to be beneficial on cultivars highly susceptible to anthracnose (e.g., Marquette, Reliance), where it can be important in blocks with a history of the disease, or in blocks where black rot and/or Phomopsis control is regularly problematic and conventional fungicides will not be used during the growing season. Otherwise, it is unlikely to be cost effective. THOROUGH coverage of the vines is essential for acceptable results. If practical, application to individual vines with a handgun or using some other system that minimizes loss of these expensive materials to non-grapevine surfaces (e.g., hooded-boom, recirculating sprayer such as the Lipco) is desirable. The low per-acre rate of * ^{NY} Sulforix is unlikely to be effective unless loss to non-target surfaces is minimal. Use of more than 15 gal/A of * ^{NY} Miller Lime Sulfur is prohibitively expensive. For NYS users: note that * ^{NY} Sulforix is only labeled for use against powdery mildew and Phomopsis and that * ^{NY} Miller Lime Sulfur is only labeled for control of Phomopsis, powdery mildew and anthracnose.
	OR * ^{NY} Sulforix	1-2 gal	

Pest(s)	Materials	Rate per Acre	Comments
5.2.9 FIRST POSTBLOOM SPRAY (10-14 days after immediate prebloom) (continued)			
Powdery mildew (continued)	OR * ^{NY} Aprovia 0.83EC	8.6-10.5 fl oz	* ^{NY} Aprovia is a Group 7 fungicide; it is a solo product (not combined with another active ingredient). * ^{NY} Aprovia has provided excellent activity against powdery mildew in multiple trials, but has not provided reliable activity against any other disease.
	OR * ^{NY} Aprovia Top 1.6EC	8.5-13.3 fl oz	* ^{NY} Aprovia Top is a relatively new product that combines the active ingredient contained in * ^{NY} Aprovia plus an effective rate of the DMI fungicide difenoconazole: both active ingredients have considerable activity against powdery mildew. CAUTION: DO NOT USE *^{NY}APROVIA TOP ON CONCORD GRAPES (CAUSES INJURY).
	OR * ^{NY} †Miravis Prime	9.2-13.4 fl oz	* ^{NY} †Miravis Prime is a combination of a new succinate dehydrogenase inhibitor fungicide (SDHI, Group 7) and fludioxonil (Group 12). In NY and PA trials, * ^{NY} †Miravis Prime has provided high levels of powdery mildew control. It is recommended that the total number of applications of Group 7 materials (* ^{NY} †Luna Experience, Pristine, Endura, and * ^{NY} Aprovia) be limited to no more than two per season, preferably not in sequence.
	OR Double Nickel LC or Double Nickel 55	2 qt 1 lb	Double Nickel is a biopesticide sold in both liquid and WDG formulations. It requires relatively short spray intervals and has provided fair to very good results in NY trials, depending on cultivar susceptibility. It has been most effective on moderately susceptible cultivars rather than highly susceptible cultivars such as Chardonnay.
	OR Oso 5SC or Ph-D 89WDG	6.5 fl oz 6.2 oz	Oso and Ph-D are relatively new products that contain the same, older active ingredient. They have provided good to very good results in NY trials when rotated with strong conventional materials. However, they should not be relied upon at this particularly critical time on <i>V. vinifera</i> and other highly susceptible cultivars.
	OR Regalia	1-4 qts (see label)	Regalia is a biopesticide that has provided fair to very good control of powdery mildew in NY trials, depending on disease pressure. It is likely to work best on less susceptible varieties or at times of year when pressure is low. Different rates are recommended depending on whether it is used alone or in a tank mix. Because Regalia requires 48 hours to activate plant defenses, applications need to start before infection. Regalia should only be considered for powdery mildew control during post bloom in an organic spray program. It should not be relied on for conventional powdery mildew control during this critical window.
	OR ProBlad Verde	18.1-45.7 fl oz	ProBlad Verde is a biopesticide that has provided fair to good control of powdery mildew in NY and PA trials. ProBlad Verde is OMRI listed and should only be considered for powdery mildew control during post bloom in an organic spray program. ProBlad Verde should not be relied on for conventional powdery mildew control during this critical window. No more than 5 applications can be made during the season, and ProBlad Verde may be applied no more than twice before alternating with a product that has a different mode of action.

6 Vineyard Weed Management

6.1 Introduction

Weeds are part of the vineyard ecosystem. Weed management decisions are based on balancing the positive and negative aspects of weed growth in the vineyard. Weeds can compete for water and nutrients, reducing vine growth; contaminate mechanically harvested fruit; provide alternate hosts for vineyard pests; and interfere with vineyard operations. Weed growth can also alter the microclimate around vines, leading to higher disease pressure and increasing the risk of spring frost. Managing weed or cover crop growth in row middles can be a powerful tool for managing overly vigorous vines, minimizing erosion, and improving equipment access in wet seasons. Weed management practices can have negative impacts on grapes if those strategies cause direct damage to the vines.

This portion of the guide primarily addresses chemical methods to control weeds in vineyards. We have attempted to include all herbicides labeled for use in grapes even though some are not commonly used in eastern United States vineyards. Not all products or use patterns are labeled for use in each state or in every region of the same state. Registrations may change, so product users should always rely on the most up to date labels for use recommendations. Herbicides are listed in the sections, “preemergence herbicides,” “postemergence herbicides,” and “herbicides for nonbearing vineyards.” Herbicides that are registered for vineyard use and may have applicability under specific circumstances are listed in the section, “specialty use herbicides.”

Cultivation is sometimes used as a weed management tool in vineyards. Low vine size restricts productivity of own-rooted *Vitis labruscana* varieties such as ‘Concord’ that generally have shallow root systems. Effective herbicide use has been shown to increase vine size and subsequent yields as compared with under-the-row cultivation, but this may be less of a concern when deep-rooted rootstocks are used, when vines are overly vigorous, or when maximum yields are not desired. Under trellis mowing and growing cover crops under the row are currently being researched in New York and Virginia. These methods might also be considered where vine growth is overly vigorous.

Cultivation and organic mulches can also be used as tools for row middle management. Excessive cultivation can lead to undesirable consequences such as soil erosion, reduced soil organic matter, and breakdown in soil structure resulting in compaction and reduced permeability. Recently cultivated soil can restrict equipment mobility needed for critical vineyard operations such as timely pesticide applications and mechanical harvest. If cultivation is used for row middle management it is suggested that negative effects be limited by not cultivating more often than necessary to suppress weed growth, to shallow (1-2")

depths only, and with the goal of reducing, rather than eliminating, weed or cover crop growth. Fall planting of ryegrass or other cover crops can be used in conjunction with cultivation to provide winter cover. Organic mulches are most effective where soil moisture and fertility are low and where low vine size restricts vineyard productivity.

6.2 Resources

Many resources for weed identification and management can be found through the Cornell *Weed Science* website at <https://cals.cornell.edu/weed-science>. Additionally, *Weeds of the Northeast* is an excellent resource covering weed identification and aspects of weed biology and ecology that relate to weed management. *Weeds of the Northeast* is available through Cornell University Press and the Lake Erie Regional Grape Program office in Portland, NY. Another valuable resource is the *Manage Weeds on your Farm* e-book available for free at www.sare.org/resources/manage-weeds-on-your-farm/. This e-book includes descriptions of many ecologically based weed management techniques as well as detailed accounts of the biology and management of the most common weed species. For a starting point in determining which herbicides are effective on individual species, the Michigan State University *Herbicide/Weed Database* (<https://herbicides.hrt.msu.edu>) may be helpful, but be sure to check that products are labeled for usage in New York grapes.

6.3 Effective Use of Herbicides

Repeated use of the same herbicides, or those with similar chemistry, can lead to a buildup of tolerant weeds, development of resistant biotypes, shifts in microbial communities that can facilitate rapid decomposition, which can reduce the length of time soil active herbicides are effective. Although herbicide options in vineyards are somewhat limited, weed management programs should not rely on the repeated use of the same herbicide modes of action. See Table 8.1 for a listing of herbicide mode of action group numbers from the Weed Science Society of America.

Table 6.3.1 Acres of vineyard sprayed at different row and spray band widths to equal one acre of land surface sprayed.

Width of Spray Band	Distance between rows		
	8 ft	9 ft	10 ft
30 inches	3.2	3.6	4.0
36 inches	2.7	3.0	3.3
42 inches	2.3	2.6	2.9
48 inches	2.0	2.3	2.5

***NY†PRINCEP Caliber 90, *NY†PRINCEP 4L**, others (simazine) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (800) 888-8372

Selected uses: Preemergence control of broadleaf weeds and early season control of annual grasses in established vineyards.

Rate: Apply 2.2-4.4 lb of 90 WDG, or 2-4 qt of 4L per acre surface sprayed. Use the lower rate on coarse-textured soils low in organic matter. Note the maximum labeled use rate has recently been reduced.

Timing: Apply to weed free soil, or use with a post-emergence herbicide, between harvest and early spring before weeds emerge. Do not use in vineyards established less than 3 years.

Comments: Princep controls a broad spectrum of annual broadleaf weeds, but season-long control of annual grasses is not usually achieved. At low soil pH, hydrolysis enhances degradation and residual activity is shortened. For season-long control of annual grasses, use a tank mix with *NY†Solicam, Prowl H₂O, oryzalin, or diuron.

PROWL H₂O (pendimethalin) – read the label

Signal word: CAUTION

Medical emergency: (800) 832-4357

Selected uses: Preemergence control of annual grasses and control or suppression of some annual broadleaf weeds in bearing and non-bearing vineyards; must be applied when vines are dormant. Note that other formulations of pendimethalin have been registered for use in non-bearing vineyards for several years, but use in bearing vineyards is restricted to the “Prowl H₂O” formulation.

Rate: Apply 3.2 to 6.3 quarts per acre surface sprayed.

Timing: Any time after fall harvest, during winter dormancy, or in the spring. Do not apply over the top of grape vines with leaves, buds, or fruit. Contact with leaves, shoots, or buds may cause injury.

Comments: The active ingredient in Prowl H₂O (pendimethalin) is chemically similar to oryzalin (both are dinitroanilines) and the spectrum of weeds they control is similar. Pendimethalin is weak in controlling some broadleaf weeds including ragweed and mustard species. Tank mix with *NY†Solicam, Chateau, simazine, diuron, or oxyfluorfen for broad-spectrum weed control.

NOTE: Prowl H₂O is also registered for use during vineyard establishment. That use is discussed in the section “Herbicides for nonbearing vineyards.”

***NY†SOLICAM DF** (norflurazon) – RESTRICTED-USE PESTICIDE IN NY – read the label

Signal word: CAUTION

Medical emergency: (866) 374-1975

Selected uses: Preemergence control of annual grasses, suppression of perennial grasses, and control or suppression of some annual broadleaf weeds in established vineyards.

Rate: Apply 1.25-5 lb per acre surface sprayed. Consult the label to determine the proper use rate, which is based on soil texture.

Timing: Apply in fall or early spring prior to weed germination. Several inches of rainfall are needed within 4 weeks of application to move the herbicide into the soil. Vines must be established at least 2 years. Some whitening maybe observed if used close to bud break.

Comments: *NY†Solicam provides excellent season-long control of annual grasses and several annual broadleaf species including velvetleaf, and will provide some suppression of yellow nutsedge, plantains, and perennial grasses. *NY†Solicam is relatively weak in controlling common annual broadleaf weeds such as pigweed and smartweed species and common lambsquarters. Apply to weed-free soil. Tank mixes with another preemergence herbicide such as Prowl H₂O, simazine, diuron, oxyfluorfen, or oryzalin are needed to attain season-long, broad-spectrum weed control. Note that *NY†Solicam is not registered for use in Nassau and Suffolk counties in New York.

SURFLAN A.S. (oryzalin) – read the label

Signal word: CAUTION

Medical emergency: (866) 673-6671

Selected uses: Preemergence control of annual grasses and control or suppression of some annual broadleaf weeds in bearing and nonbearing vineyards.

Rate: Apply 2-6 qt per acre surface sprayed. Length of control depends on the rate applied. The total amount allowed per year is 12 qt per acre surface sprayed with a minimum of 2.5 months between applications.

Timing: Apply in the fall or spring prior to weed germination. Do not apply to newly planted vines until soil has settled.

Comments: Oryzalin may be applied safely to coarse-textured, low organic matter soils where other residual herbicide options are limited. It is not recommended for use on soils with an organic matter content of greater than 5 percent. Apply to weed free soil; residues on the soil surface can also reduce its effectiveness. Oryzalin is very weak in controlling some broadleaf weeds including ragweed and mustard species. Tank mix with *NY†Solicam, diuron, simazine, or oxyfluorfen for broad-spectrum season-long weed control. Note: the product “Surflan A.S. Specialty Herbicide” is only registered for use in non-bearing vineyards.

NOTE: Oryzalin is also registered for use during vineyard establishment. That use is discussed in the section “Herbicides for nonbearing vineyards.”

6.5 Postemergence Herbicides

AIM (carfentrazone-ethyl) – read the label

Signal word: CAUTION

Medical emergency: (800) 331-3148

Selected uses: Postemergence control of certain susceptible broadleaf weeds and burn down of grapevine suckers

Rate: For broadleaf weed control, apply up to 2 fl. oz. per acre surface sprayed. Lower rates can be used to control small seedling weeds at the 2 to 3-leaf stage; higher rates are needed for larger weeds up to the 6-leaf stage. Applications to weeds beyond the six-leaf stage may result in only partial control. For burn down of grapevine suckers, apply at the maximum use rate (2 fl. oz.) per acre surface sprayed. Add a non-ionic surfactant (NIS) containing at least 80% active ingredient at 2 pt. per 100 gallons, or a crop oil concentrate (COC) at one gallon COC per 100 gallons, or methylated seed oil (MSO).

Timing: Aim may be applied at any time during the season, but do not allow spray mist to contact desirable fruit, foliage, or green bark. Suckers and other undesirable growth must be treated when the tissue is young (not mature or hardened off). Multiple applications per season are allowed, but do not apply more than 7.9 fl. oz. per season. Do not make applications less than 14 days apart or within 3 days of harvest.

Comments: Aim is very effective in controlling grapevine suckers, and also controls some small broadleaf weeds. Treated suckers turn brown within 1-2 days of application, but multiple applications may be necessary to obtain season-long sucker control. Refer to the label for a list of susceptible broadleaf weeds. Aim may be tank-mixed with other preemergence and postemergence herbicides; observe the other product's label restrictions.

^*GRAMOXONE SL 2.0 (paraquat) – RESTRICTED-USE PESTICIDE – read the label

Signal word: DANGER

Medical emergency: (800) 888-8372

Selected uses: Postemergence burn down of all weeds in new or established vineyards and burn down of grapevine suckers up to 8 inches long.

Rate: See labels as multiple products are registered. ^*Gramoxone SL 2.0 contains 2 lb. of the active ingredient, paraquat, per gallon. Apply 2.5-4 pt per acre surface sprayed. Always add a nonionic surfactant (NIS) or crop oil concentrate (COC). Add NIS at 1 pt per 100 gal (75% or more surface-active agent), or NIS at 2 pt per 100 gal (50-74% surface-active agent), or COC at 1 gal per 100 gal. The label permits applications in as low as 10 gal per treated acre, but spray volume should be increased as necessary to obtain complete coverage of

target weeds or suckers without runoff from the target foliage.

Timing: *Gramoxone should be applied to emerged weeds when they are small and succulent. Weeds 1-6 inches tall are easiest to control. Contacted plant foliage wilts and desiccates within hours of application, with complete necrosis in 1-3 days. For burndown of grapevine suckers, treat when sucker growth is no more than 8 inches long. For mature woody weeds, perennial weeds, late-germinating weeds, and green suckers, retreatment or spot -treatment may be necessary. Late season applications to weeds should be made to avoid contact with desirable foliage. Contact of spray or mist with fruit is expressly prohibited on the label and is not a legal use of the product. *Gramoxone applications are most effective under cool, cloudy conditions when drying time is slow and uptake into target foliage is optimal.

Comments: *Gramoxone is a dangerous poison and can be fatal if swallowed or inhaled. It is harmful if absorbed through the skin and can cause substantial eye injury. Follow all label precautions when mixing and applying. In vineyards where chemical suckering is undesirable, avoid contact with suckers by making a directed application under appropriate environmental conditions or use shielded equipment. Tank mixes with appropriate preemergence herbicides are recommended for broad-spectrum control and residual activity. Note that newer paraquat labels require: use only by certified applicators, the certified applicator to complete an EPA-approved paraquat training every 3 years, and use of closed transfer systems when using smaller containers. Growers using paraquat with older, still registered labels, can use the product according to the label on those containers.

MATRIX FNV (rimsulfuron) – read the label

Signal word: CAUTION

Medical emergency: (800) 441-3637

Selected uses: Primarily a pre-emergence product but will control many annual grasses and broadleaf weeds while they are small. Refer to Section 6.4 for additional information.

Comments: A FIFRA Section 2(ee) recommendation for unlabeled pest in New York State allowing to be used post-emergence to target hedge bindweed. 2(ee) recommendations for this use also available for the rimsulfuron-containing products Matrix FNV, Matrix SG, Pruvion, Grapple, Hinge, Solida, and Rimgro. Users must have a copy of the 2(ee) recommendation in their possession at the time of use. Ineffective on field bindweed. Refer to the recent Cornell identification guide for more details (<https://hdl.handle.net/1813/70190>).

POAST (sethoxydim) – read the label

Signal word: WARNING

Medical emergency: (800) 832-4357

7 Sprayer Technology

7.1 Solutions for Safer Spraying – Engineering Controls

Keeping pesticide exposure to a minimum should be a chief concern of any pesticide applicator. To reduce the risks associated with handling and applying pesticides, devices known as **engineering controls** can be used to reduce or practically eliminate exposure to hazardous chemicals.

7.1.1 Areas of Exposure

1. Loading the Sprayer

Closed transfer systems. Closed transfer systems allow concentrated pesticide to be moved from the original shipping container to the sprayer mix tank with minimal or no applicator contact. Many systems provide a method to measure the concentrated pesticide. Some systems also include a container rinsing system. Currently available closed transfer systems use a probe inserted into the pesticide container, a connector on the container that mates to a similar connector on the application equipment, or a vacuum-type (venturi) system that uses flowing water to transfer the chemical from the container.

Induction bowls. Induction bowls are metal, plastic or fiberglass hoppers attached to the side of the sprayer or the nurse tank that allow pesticides to be added to the mix tank without the applicator climbing onto the spray rig. Pesticides are poured into the bowl and water is added to flush out the bowl and carry the pesticide to the spray tank. Often a rinse nozzle is mounted inside the bowl for rinsing out empty pesticide containers. Typically, induction bowls are raised out of the way during spraying and lowered to about 3 feet above ground when the sprayer is being loaded.

Direct pesticide injection system. Direct pesticide injection systems allow pesticides to be mixed directly with water in the sprayer plumbing system rather than in the main spray tank. The pesticide is pumped from its container and mixed with the water either in a manifold or at the main water pump. Only clean water is held in the main tank of the sprayer. An electronic controller and up to four pumps adjust the amount of concentrated pesticide that is injected into the water stream, allowing for variable application rates.

Container rinse system. Container rinse systems consist of a rinse nozzle and a catch bowl that traps the container washings (rinsate). The empty container is placed over the rinse nozzle and a jet of water cleans the inside of the container. The rinsate caught in the bowl is pumped into the spray tank to be used along with the spray mixture. Often rinse nozzles are installed in chemical induction bowls. Most closed transfer systems also provide a way of rinsing containers and piping the rinse water into the spray tank.

2. Reducing Exposure at the Boom

Boom folding/extending. Manually folding booms can be a source of operator contamination because the boom can be covered with pesticide from drift or dripping nozzles. Consider the use of hydraulic or mechanical folding methods.

Diaphragm check valves. Typically, when a sprayer is shut off and as the system pressure drops, any liquid remaining in the boom piping drips from the nozzles, possibly dripping onto the boom or even the operator. Diaphragm check valves installed at each nozzle prevent this by using a spring-loaded rubber diaphragm to close off the flow of liquid once the system pressure drops below about 10 pounds per square inch. When the sprayer is switched on and system pressure builds up, the valve opens and allows the liquid to flow through the nozzles.

Multiple nozzle bodies. Contamination can occur when operators change or unclog nozzles during an application. Multiple nozzle bodies (or turret nozzles) allow operators to switch between nozzles with a turn of the nozzle body rather than having to unscrew or undo a threaded or bayonet fitting.

Hand wash water supply. Providing adequate wash water is essential (and often required). A simple container with a hand-operated valve can be mounted on the side of the sprayer to provide clean water for hand washing and personal hygiene.

3. Drift and Contaminated Clothing in Cabs

Cab filtration using carbon filters. Carbon filtration systems are used to remove pesticide odor and pesticide-laden mist from fresh air used in a tractor or self-propelled sprayer cab. Carbon filtration systems are often a standard feature on self-propelled sprayers. Many factory installed tractor cabs offer optional filtration systems.

Protective clothing lockers. To prevent contamination of the tractor or sprayer cab interior, protective clothing should be removed before entering the cab. A few sprayer companies offer a simple compartment (or locker) mounted to the side or front of the sprayer where protective clothing can be stored.

4. Controlling Drift

Low-drift nozzles. Low-drift nozzles create larger-size droplets than conventional nozzles. The larger droplet sizes are less prone to drift, reducing environmental and operator contamination.

Air induction (twin fluid) nozzles. These nozzles allow air to mix with the spray liquid, creating large, air-filled droplets with virtually no fine, drift-prone droplets. The droplets explode when they contact their target and offer similar coverage to droplets from conventional, finer-spray nozzles.

selecting and changing nozzles" or "Calibration of airblast sprayers for orchards part 2 measuring liquid flow".

7.3.1 Travel Speed Calibration

Sprayer travel speed influences spray deposition and is a critical factor in maintaining accurate application rates. Studies to determine the effect of travel speed on average spray deposition agree that the higher the travel speed, the greater the variability in spray deposit. This variability is an important factor where uniformity of spray coverage throughout the canopy is required. Conclusions from research were drawn using travel speeds of 1-4 mph.

Factors affecting travel speed include:

- weight of sprayer to be pulled.
- slope of terrain.
- ground conditions traveled over (wheel slippage).

The best way to measure travel speed is to pull a sprayer half full of water over the same type of terrain on which the actual sprayer will be operated.

Using a tape measure, set up a test course at least 100 feet long. Do not pace the distance. The longer the course, the smaller the margin of error.

Run the course in both directions. Use an accurate stopwatch to check the time required to travel the course in each direction. Average the two runs and use the following to calculate the speed in MPH.

Formula $\frac{\text{Ft. traveled}}{\text{Sec. traveled}} \times \frac{60}{88} = \text{MPH}$

Your figures:

Tractor gear Engine revs.

$$\frac{\text{ft.}}{\text{sec}} \times \frac{60}{88} = \underline{\hspace{1cm}} \text{MPH}$$

An alternative to using the above method is to purchase a hand-held GPS receiver. A number of systems are available from electronics stores, hunting equipment suppliers, and online. Costing ranges from \$80-150. The small device is portable so can be used in all tractors to determine forward speed in specific tractor gears at known engine RPM.

7.3.2 Airblast Sprayer Calibration

- use clean water

1. Pressure check

Place the pressure gauge on the nozzle fitting farthest away from the pump and turn the sprayer on. If pressure is lower at the nozzle than specified, increase pressure at the regulator.

Pressure at nozzle _____ psi

Pressure at sprayer gauge psi

2. Nozzle output

Use a flow meter (obtainable from Gemplers, Spraying Systems, or other sources) attached to individual nozzles OR connect hoses to each of the nozzles and measure the flow from each nozzle into a calibrated jug. Record and total your results using Figure 7.3.1.

Calculate gallons per acre using the following formula.

Formula: $\frac{\text{Total GPM X 495}}{\text{mph X row spacing (ft.)}} = \text{GPA}$

Your figures: $\frac{\text{GPM X 495}}{\text{mph X ft.}} = \text{GPA}$

Replace all nozzle tips which are more than 5% inaccurate.

[illegible]

Figure 7.3.1. Airblast Sprayer Calibration

7.10 Going Spraying!

Mixing Procedures



Safety and the Law

- Always remain alert; pesticides are potentially dangerous to the operator and the environment.
- Tractors and sprayers are dangerous machines and care should be taken when operating them.
- Always follow federal and state laws concerning applicator certification and handling, application and disposal of pesticides.

Always read the label for detailed application information and keep a record.

WORK REPORT No.	
INSTRUCTIONS	
DATE	
FARM & FIELD	
CROP	
AREA	
OPERATION	
Spraying	
PRODUCT	
QUANTITY	
Rate	
Comments	
OPERATOR REPORT	
NAME	
DATE	
MACHINE USED	
TIME START	
TIME STOP	
ACUALLY PRODUCT USED	
TOTAL	
TIME	
FUEL	
OIL	
WATER	
PUMP	
COMMENTS	

The seven Ps of machinery management.

Proper prior planning prevents poor performance.



- Fill the tank on level ground per label instructions. If none are given, fill the tank half full with clean water.



- Prime the pump with water, if needed.

8 Pesticides for New York and Pennsylvania Vineyards

8.1 Herbicides

Common Name	Trade Name	Formulation	WSSA Group Number (Resistance Management)	Days to Harvest	Restricted Entry Interval (REI)	EPA Reg. Number
carfentrazone-ethyl	Aim EC	EC	14	3	12	279-3241
clethodim	^Select 2EC	2EC	1	1 year	24 hr	59639-3-1381
	^Volunteer	2 EC	1	1 year	24 hr	59639-3-55467
	* ^{NY} Select Max	0.97 EC	1	1 year	24 hr	59639-132
dichlobenil	Casoron 4G	4G	20	0	24 hr	70506-519
diuron	Direx 4L	4L	7	0	12 hr	66222-54
	Karmex DF	80 DF	7	0	12 hr	66222-51
fluazifop-P-butyl	†Fusilade DX	2EC	1A	50	12 hr	100-1070
flumioxazin	Chateau SW	51 WDG	14	60	12 hr	59639-99
glufosinate-ammonium	†Rely 280	2.34 EC	10	14	12 hr	7969-448
glyphosate	Multiple	Multiple	9	See label	See label	Multiple
indaziflam	* ^{NY} †Alion	SC	29	14	12 hr	264-1106
isoxaben	^Gallery	75DF	21	1 year	12 hr	62719-145
napropamide	Devrinol DF-XT	50DF	15	70	24 hr	70506-36
norflurazon	* ^{NY} †Solicam	80DF	12	60	12 hr	61842-41
oryzalin	Surflan	4AS	3	0	24 hr	70506-43
oxyfluorfen	Goal 2XL	2EC	14	^a	24 hr	62719-424
	GoalTender	4 EC	14	^a	24 hr	62719-447
paraquat	^*Gramoxone SL 2.0	2L	22	0	24 hr	100-1431
pelargonic acid	Scythe	4.2EC	unclassified	0	12 hr	10163-325
pendimethalin	Prowl	3.3EC	3	1 year	24 hr	241-337
	Prowl H ₂ O	3.8EC	3	21	24 hr	241-418
rimsulfuron	Matrix FNV	25DF	2	14	4 hr	352-671
sethoxydim	Poast	1.5EC	1	50	12 hr	7969-58
simazine	* ^{NY} †Princep Caliber 90 and others	90WDG	5	0	12 hr	100-603
	* ^{NY} †Princep 4L and others	4L	5	0	12 hr	100-526
trifluralin	Multiple	various	3	^b	See label	Multiple
trifluralin + isoxaben^b	^Snapshot	2.5TG	3, 21	1 year	12 hr	62719-175

Notes: ^a Do not apply after bud swell

^b Apply and incorporate before planting

* Federal restricted use pesticide.

*^{NY} Restricted use pesticide in New York State

† Not for use in Nassau/Suffolk Counties in New York

^ Not registered for use in New York State at press time

8.2 Insecticides and acaricides (continued)

Common Name	Trade Name	Formulation	Days to Harvest	Restricted Entry Interval (REI)	EPA Reg. Number
imidacloprid	* ^{NY} Prey	1.6F	0	12 hr	34704-894
	* ^{NY} Admire Pro	4.6L	30 (soil); 0 (foliar)	12 hr	264-827
	* ^{NY} Alias 4F	4F	30 (soil); 0 (foliar)	12 hr	66222-156
imidacloprid + beta-cyfluthrin	*Leverage 360		3	12 hr	264-1104
malathion	Drexel Malathion 5EC	5 EC	3	24 hr	19713-217
	Malathion 8 Aquamul	8 EC	3	24 hr ¹	34704-474
methoxyfenozide	* ^{NY} †Intrepid 2F	2F	30	4 hr	62719-442 (SLN NY-190006 in NYS)
methoxyfenozide + spinetoram	^Intrepid Edge	LF	30	4 hr	62719-666
phosmet	* ^{NY} Imidan	70W	7 or 14 (rate based - see label)	14 days ¹	10163-169
potassium salts of fatty acids	M-Pede	insecticidal soap	0	12 hr	10163-324
pyrethrins	Evergreen	EC	0 (when dry)	12 hr	1021-1770
	Pest-Kote	EC	0 (when dry)	12 hr	92035-1-97839
pyrethrins + piperonyl butoxide	*Pyronyl Crop Spray	EC	0 (when dry)	12 hr	89459-26
pyridaben	†Nexter	SC	7	12 hr	81880-28-10163
pyriproxyfen	Knack	EC	21	12 hr	59639-95
spinetoram	Delegate	WG	3	4 hr	62719-541
spinosad	Entrust	WP	7	4 hr	62719-282
	Entrust SC	SC	3	4 hr	62719-621
spirotetramat	Movento		7	24 hr	264-1050
spirotetramat + pyriproxyfen	Senstar	SE	21	24 hr	59639-243
thiamethoxam	* ^{NY} †Actara	WDG	5	12 hr	100-938
thiamethoxam + chlorantraniliprole	* ^{NY} †Voliam Flexi	WG	14	12 hr	100-1319
zeta-cypermethrin	*Mustang Maxx		1	12 hr	279-3426
zeta-cypermethrin + bifenthrin	*Hero	EC	30	12 hr	279-3315

* Federal restricted-use pesticide

*^{NY} Restricted-use pesticide in New York State

† Not for use in Nassau/Suffolk Counties in New York

^ Not registered for use in New York State at press time

¹ Multiple REIs – see label for details**8.3 Fungicides**

Common Name	Trade Name	Formulation	Days to Harvest	Restricted Entry Interval (REI)	EPA Reg. Number
ametoctradin + dimethomorph	* ^{NY} †Zampro	4.4SC	14	12 hr	7969-302
<i>Aureobasidium pullulans</i> strains DSM 14940 and 14941	Botector	80WP	UDH ²	4 hr	86174-3
azoxystrobin	Abound	2SC	14	4 hr	100-1098

Tips for Laundering Pesticide-Contaminated Clothing

Pre-Laundering Information

Remove contaminated clothing **before** entering enclosed tractor cabs.

Remove contaminated clothing **outdoors** or in an entry. If a granular pesticide was used, shake clothing outdoors. **Empty pockets and cuffs.**

Save clothing worn while handling pesticides for that use only. Keep separate from other clothing **before, during, and after** laundering.

Wash contaminated clothing after **each** use. When applying pesticides daily, wash clothing **daily**.

Clean gloves, aprons, boots, rigid hats, respirators, and eyewear by scrubbing with detergent and warm water. Rinse thoroughly and hang in a clean area to dry.

Take these **precautions** when handling contaminated clothing:

- Ventilate area.
- Avoid inhaling steam from washer or dryer.
- Wash hands thoroughly.
- Consider wearing chemical-resistant gloves.
- Keep out of reach of children and pets.

Air

Hang garments outdoors to air.

Pre-rinse

Use one of three methods:

1. Hose off garments outdoors.
2. Rinse in separate tub or pail.
3. Rinse in automatic washer at full water level.

Pretreat (heavily soiled garments)

Use heavy-duty liquid detergent.

Washer Load

Wash garments separate from family wash.

Wash garments contaminated with the same pesticide together.

Never use the “sudsaver” feature on your machine when laundering pesticide-soiled clothes.

Load Size

Wash only a few garments at once.

Water Level

Use full water level.

Water Temperature

Use **hot** water, as hot as possible.

Wash Cycle

Use **regular** wash cycle, at least 12-minutes.

Laundry Detergent

Use a **heavy-duty** detergent.

Use amount recommended on package or more for heavy soil or hard water.

Remember to use high-efficiency (HE) detergents in HE and front-loading washers.

Rinse

Use a **full** warm rinse.

Rewash

Rewash contaminated garments **two or three times** before reuse for more complete pesticide removal.

Dry

Line drying is preferable to avoid contaminating dryer.

Clean Washer

Run complete, but empty, cycle.
Use **hot water and detergent**.

PESTICIDE EMERGENCY NUMBERS

Emergency responder information on pesticide spills and accidents...

CHEMTREC800-424-9300

For pesticide information...

National Pesticide Information Center800-858-7378

To Report Oil and Hazardous Material Spills in New York State...

NYS Spill Hotline800-457-7362

Poison Control Centers

Poison Control Centers nationwide800-222-1222

To order additional copies of this publication, contact your local Cornell Cooperative Extension office or order directly from the Cornell Store, Attn: PMEP, 135 Ho Plaza, Cornell University, Ithaca, NY 14853. Phone 844-688-7620. Additional copies and online access can also be ordered online at: <https://store.cornell.edu/books/cornell-cooperative-ext-pmep-guidelines>.

This publication is issued to further Cooperative Extension work mandated by acts of Congress of May 8 and June 30, 1914. It was produced with the cooperation of the U.S. Department of Agriculture, Cornell Cooperative Extension, New York State College of Agriculture and Life Sciences, New York State College of Human Ecology, and New York State College of Veterinary Medicine, at Cornell University. Cornell Cooperative Extension provides equal program and employment opportunities.

Alternative formats of this publication are available on request to persons with disabilities who cannot use the printed format. For information call or write the Office of the Director, Cornell Cooperative Extension, 365 Roberts Hall, Ithaca, NY 14853 (607-255-2237).

This information is presented with the understanding that no product discrimination is intended and no endorsement of any product mentioned or criticism of unnamed products is implied.

Published by:
Cornell Integrated Pest Management Program
Pesticide Safety Education
CALS Surge Facility
525 Tower Road
Cornell University
Ithaca, New York 14853-7401
607.255.1866
Michael Helms, Managing Editor (mjh14@cornell.edu)

© 2025 Cornell University. All rights reserved.